

ON TORSION-FREE GROUPS WITH FINITE REGULAR FILE BASES

ALEXEY MURANOV

ABSTRACT. The following question was asked by V. V. Bludov in The Kurovka Notebook in 1995: If a torsion-free group G has a finite system of generators $a_1, \dots, a_n$ such that every element of G has a unique presentation in the form $a_1^{k_1} \dots a_n^{k_n}$ where $k_i \in \mathbb{Z}$, is it true that G is virtually polycyclic? The answer is “not always.” A counterexample is constructed in this paper as a group presented by generators and defining relations.

CONTENTS

1. Introduction	1
2. Group construction	3
3. Definitions	3
4. The Main Lemma	6
5. Properties of the group	8
6. Comments	38
Acknowledgements	39
References	39

1. INTRODUCTION

Definition. A group is *polycyclic* if it has a finite subnormal series with cyclic factors. A group is *virtually polycyclic* if it has a polycyclic subgroup of finite index.

Subgroups and homomorphic images of polycyclic and virtually polycyclic groups are polycyclic or, respectively, virtually polycyclic. Finitely generated abelian groups are polycyclic. Free groups of rank at least 2 are not virtually polycyclic.

Vasiliy Bludov posed the following question in 1995 (Problem 13.11 in [MK95b]; see also Question 6 in [Blu95b]):

Date: February 1, 2008.

2000 Mathematics Subject Classification. Primary 20F05; Secondary 20F06.

Key words and phrases. Group presentation, van Kampen’s lemma, diagram with selection, S-diagram, virtually polycyclic group, boundedly generated group, file basis.

This work was supported in part by the NSF grant DMS 0245600 of Alexander Ol’shanskiy and Mark Sapir.

If a torsion-free group G has a finite system of generators $a_1, \dots, a_n$ such that every element of G has a unique presentation in the form $a_1^{k_1} \dots a_n^{k_n}$ where $k_i \in \mathbb{Z}$, is it true then that G is virtually polycyclic?

The following theorem answers Bludov's question negatively:

Theorem. *Provided $n \geq 63$, there exists a group G and pairwise distinct elements $a_1, \dots, a_n$ in G such that:*

- (0) G is generated by $\{a_1, \dots, a_n\}$;
- (1) every $n - 21$ elements out of $\{a_1, \dots, a_n\}$ freely generate a free subgroup such that every two elements of this subgroup F are conjugate in G only if they are conjugate in F itself (in particular, G is not virtually polycyclic);
- (2) for every element g of G , there is a unique n -tuple $(k_1, \dots, k_n) \in \mathbb{Z}^n$ such that $g = a_1^{k_1} \dots a_n^{k_n}$;
- (3) G is torsion-free;
- (4) G is the direct limit of a sequence of hyperbolic groups with respect to a family of surjective homomorphisms;
- (5) G is recursively presented and has solvable word and conjugacy problems.

(The *direct limit* in condition (4) is the direct limit in the sense of Bourbaki; see “Algebra I.”)

This theorem is proved in the end of Section 5 of this paper.

In terms of *file bases* (see definitions in [Blu95b]), item (2) of the theorem states that the elements $a_1, \dots, a_n$ in this order form a *regular* file basis of G . The argument below shows that every group with a 2-element file basis is polycyclic (therefore, the assumption “ $n \geq 63$ ” cannot be replaced with “ $n \geq 2$ ”).

Every group that has an n -element file basis is *n -boundedly generated* (i.e., is the product of an n -tuple of its cyclic subgroups). The following two theorems (see [Itô55] and [LR80]) together yield that every 2-boundedly generated group is polycyclic:

Theorem (Itô, 1955). *If a group G has abelian subgroups A and B such that $G = AB$, then G is metabelian.*

Theorem (Lennox, Roseblade, 1980). *If a soluble group G has polycyclic subgroups A and B such that $G = AB$, then G is polycyclic.*

Some examples of non-polycyclic groups with finite regular file bases and with torsion are known. It is shown in [Blu95b] (see Example 3 therein) that the semidirect product of the rank- n free abelian group $\mathbb{Z}^n$ with the symmetric group S_n defined via the action of S_n on $\mathbb{Z}^n$ by naturally permuting the standard generators has an n -element regular file basis of infinite-order elements. If $n \geq 5$, then such a semidirect product is virtually polycyclic but not polycyclic.

The primary tool used to establish properties of the group (or rather the series of groups) constructed in this paper is the Main Theorem in [Mur05].

2. GROUP CONSTRUCTION

Take an arbitrary integer $n \geq 63$. Choose a positive $\lambda_1 < 1$ such that

$$(2.1) \quad \left(4 + \frac{2n\lambda_1}{1 - \lambda_1}\right) \lambda_1 \leq \frac{1}{n}.$$

(It suffices if, for example, $0 < \lambda_1 \leq 1/(5n)$.)

Let $\mathfrak{A} = \{x_1, \dots, x_n\}$ be an n -letter alphabet. Call a group word w over $\mathfrak{A}$ *regular* if it has the form $x_1^{k_1} x_2^{k_2} \dots x_n^{k_n}$ where $k_i \in \mathbb{Z}$. (In particular, regular group words are cyclically reduced.) Call a group word w *counter-regular* if w^{-1} is regular. The empty word is both regular and counter-regular. So is every group word that is a *letter power*.

Impose an order on the set of all reduced group words in the alphabet $\mathfrak{A}$ to make it order-isomorphic with the set of natural numbers. For example, use the *deg-lex* order: $x_1, x_1^{-1}, x_2, \dots, x_n^{-1}, x_1^2, x_1 x_2, x_1 x_2^{-1}, \dots$. Define a sequence of sets $\{\mathcal{R}_i\}_{i=0}^{+\infty}$ inductively.

First, let $\mathcal{R}_0 = \emptyset$.

Second, if $i > 0$ and every group word over $\mathfrak{A}$ equals a regular word modulo the relations $r = 1, r \in \mathcal{R}_{i-1}$, then let $\mathcal{R}_i = \mathcal{R}_{i-1}$.

Last, if $i > 0$ and some group word over $\mathfrak{A}$ is not equal to any regular word modulo the relations $r = 1, r \in \mathcal{R}_{i-1}$, then let w_i be the least reduced group word (with respect to the chosen order) that does not start with $x_1^{\pm 1}$, does not end with $x_n^{\pm 1}$, and is not equal to any regular word modulo the relations $r = 1, r \in \mathcal{R}_{i-1}$ (clearly such w_i exists). Choose a natural number m_i such that

$$(2.2) \quad i > j \Rightarrow m_i \neq m_j,$$

$$(2.3) \quad i > j \Rightarrow nm_i + |w_i| \geq nm_j + |w_j|,$$

$$(2.4) \quad \lambda_1(nm_i + |w_i|) \geq |w_i|.$$

(These conditions may be satisfied by choosing a sufficiently large m_i .) Define

$$(2.5) \quad r_i = x_1^{m_i} x_2^{m_i} \dots x_n^{m_i} w_i^{-1},$$

and let $\mathcal{R}_i = \mathcal{R}_{i-1} \cup \{r_i\}$. Note that $|r_i| = nm_i + |w_i|$.

Eventually, let

$$(2.6) \quad \mathcal{R} = \bigcup_{i=1}^{+\infty} \mathcal{R}_i,$$

and let G be the group defined by the presentation $\langle \mathfrak{A} \mid r = 1, r \in \mathcal{R} \rangle$.

Observe that all elements of $\mathcal{R}$ are cyclically reduced.

Along with $\{\mathcal{R}_i\}_{i=0}^{+\infty}$, the sequences $\{w_i\}_{i=1, \dots}$, $\{m_i\}_{i=1, \dots}$, and $\{r_i\}_{i=1, \dots}$, infinite or finite, have been constructed.

Note that the above construction scheme does not specify $\mathcal{R}$ uniquely and is flexible as to the choice of $\{w_i\}_{i=1, \dots}$ and $\{m_i\}_{i=1, \dots}$.

3. DEFINITIONS

Most definitions needed for reading this paper may be found in [Mur05]. Standard terms are also defined in [LS01], [CCH81], [Ol'91] in similar ways.

For instance, geometric definitions of *maps* and *diagrams* are given in [Ol'91], and combinatorial ones in [Mur05]. Some definitions are reviewed in this section.

If X is a set, then $\|X\|$ denotes the cardinality of X .

A *graph* Γ is an ordered pair $(\Gamma(0), \Gamma(1))$ where $\Gamma(0)$ is the set of *vertices* of Γ , and $\Gamma(1)$ is its set of *edges*. It will always be assumed that $\Gamma(0)$ is nonempty. In the context of this paper, *graph* is a synonym of *combinatorial 1-complex*. A *combinatorial 2-complex* Φ is an ordered triple $(\Phi(0), \Phi(1), \Phi(2))$ where $\Phi(0)$, $\Phi(1)$, and $\Phi(2)$ are its sets of vertices, edges, and *faces*, respectively.

If v is a vertex of a complex Γ , then the number of all edges of Γ incident to v plus the number of all *loops* incident to v (thus loops are counted twice) is called the *degree* of the vertex v and is denoted by $d(v)$ or $d_\Gamma(v)$.

Here follow several definitions related to *paths* in combinatorial complexes.

The length of a path p is denoted by $|p|$. A *cyclic path* is a path whose *terminal* and *initial* vertices coincide. A *cycle* is the set of all *cyclic shifts* of a cyclic path. The cycle represented by a cyclic path p shall be denoted by $\langle p \rangle$. The *length* of a cycle c , denoted by $|c|$, is the length of an arbitrary representative of c . A *trivial cycle* is a cycle of length zero. A path p is a *subpath of a cycle* c if for some representative r of c and for some natural n , p is a subpath of r^n (i.e., of the product of n copies of r).

A path is *reduced* if it does not have a subpath of the form ee^{-1} where e is an oriented edge. A cyclic path is *cyclically reduced* if it is reduced, and its first oriented edge is not inverse to its last oriented edge. (For example, all trivial paths are cyclically reduced.) A cycle is *reduced* if it consists of cyclically reduced cyclic paths. A path is *simple* if it is nontrivial, reduced, and none of its *intermediate* vertices appears in it more than once. A cycle is *simple* if it consists of simple cyclic paths. A path is *maximal* in some set of paths if it is not a proper subpath of any other path in this set.

An *oriented arc* in a complex Γ is a simple path whose all intermediate vertices have degree 2 in Γ . A *non-oriented arc*, or simply an *arc*, is defined as a pair of mutually inverse oriented arcs. Edges and oriented edges shall be viewed as special cases of arcs and oriented arcs, respectively. Two arcs, or oriented arcs, are said to *overlap* if they have common edges. A set of arcs, or oriented arcs, is called *non-overlapping* if no two distinct elements of this set overlap. An arc u *lies* on a path p if at least one of the oriented arcs associated with u is a subpath of p .

The remaining definitions concern maps and diagrams.

Every oriented face of a 2-complex has a unique *boundary cycle*. Every (non-oriented) face of an oriented combinatorial surface has a unique *contour*, which is defined as the boundary cycle of the corresponding oriented face. The contour of a face Π shall be denoted by $\partial\Pi$. Thus the formula $\langle p \rangle = \partial\Pi$ means that the path p is a representative of the contour of the face Π . The *degree* of a face Π is the length of the contour of Π , denoted by $|\partial\Pi|$.

A *map* is a connected subcomplex of an oriented combinatorial sphere together with a set of cycles called its *contours* (see [Mur05] for a precise definition). In every map, each oriented edge appears either in the contour

of some face of the map or in a contour of the map. The union of all contours of a map Δ shall be denoted by $\partial\Delta$. A map without faces is called *degenerate*. A map is *simple* if all of its contours are simple cycles without common vertices. A *disc map* is a map with exactly one contour. A disc map is simple if and only if its underlying 2-complex is a *combinatorial disc*. An *annular map* is a map with exactly two contours.

The contours of faces in a disc map shall be thought of as *oriented counter-clockwise*, and the contour of the map itself shall be thought of as *oriented clockwise* (this is consistent with the way disc maps are pictured in this paper).

A *selection* on a face Π of a map Δ is a set of nontrivial reduced subpaths of the contour of Π closed under taking nontrivial subpaths. A *selection* on a map Δ is a set of nontrivial reduced subpaths of the contours of faces of Δ closed under taking nontrivial subpaths. An *S-map* is a map with selection. A path in an S-map is *selected* if it belongs to the selection. A path p is *double-selected* if both p and p^{-1} are selected. An arc is *double-selected* if both corresponding oriented arcs are selected. An external arc is *selected* if one of the corresponding oriented arcs is selected.

A *diagram* over a presentation $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$ is a map together with a function which labels every oriented edge of this map with a letter from $\mathfrak{B}^{\pm 1}$ so that mutually inverse oriented edges are labelled with mutually inverse group letters, and the contour of every face has a representative “labelled” with an element of $\mathcal{S}^{\pm 1}$. A diagram together with a selection on the underlying map is called an *S-diagram*.

In a diagram over $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$, let the *label* of an oriented edge be the element of $\mathfrak{B}^{\pm 1}$ that labels this oriented edge, and let the *label* of a path be the group word that “reads” on this path. Let the *label* of a (non-oriented) edge be the letter x of $\mathfrak{B}$ such that x^{+1} and x^{-1} label the two corresponding oriented edges. If q is an oriented edge or a path, the label q shall be denoted by $\ell(q)$.

Sometimes it is convenient to distinguish between a letter x of the alphabet $\mathfrak{B}$ and the corresponding letter of the group alphabet $\mathfrak{B}^{\pm 1}$ denoted by x^{+1} or simply by x . Call letters of the alphabet $\mathfrak{B}$ *basic letters* and letters of the group alphabet $\mathfrak{B}^{\pm 1}$ *group letters*. Thus, the basic letter x_1 , the group letter x_1 , and the one-letter group word x_1 are three different things. The labels of edges of a diagram are basic letters, but the labels of oriented edges are group letters. Say that a group word w has a basic letter x in it if at least one of the two group letters $x^{\pm 1}$ occurs in w .

The *mirror copy* of a map or diagram is the map or diagram, respectively, with the same underlying 2-complex, and the same labelling function in the case of diagram, but with the opposite orientation of all the faces and contours.

According to van Kampen’s lemma (see [Ol’91]), a relation $w = 1$ is a consequence of a set of relations $\{r = 1 \mid r \in \mathcal{S}\}$ if and only if there exists a disc diagram Δ over $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$, where $\mathfrak{B}$ is an appropriate alphabet, such that the label of some representative of $\partial\Delta$ is w . Such a diagram is called a *deduction diagram* for the relation $w = 1$.

Call a pair of distinct faces $\{\Pi_1, \Pi_2\}$ in a diagram Δ *immediately cancellable* if there exist paths p_1 and p_2 with a nontrivial common initial subpath such that $\langle p_1 \rangle = \partial\Pi_1$, $\langle p_2^{-1} \rangle = \partial\Pi_2$, and $\ell(p_1) = \ell(p_2)$. A diagram without immediately cancellable pairs of faces is called *weakly reduced*. (In [Mur05] immediately cancellable pairs of faces and weakly reduced diagrams are called simply *cancellable* and *reduced*, respectively.) If there exists a deduction diagram for a relation $w = 1$ over a presentation $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$, then such a diagram may be chosen weakly reduced, and even reduced in the sense of [Ol'91], which is a stronger property (take an arbitrary deduction diagram and “eliminate” immediately cancellable pairs).

Call a group presentation $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$ *strongly aspherical* if no spherical diagram over $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$ is weakly reduced.

4. THE MAIN LEMMA

This paper relies on a technical result of [Mur05] called there “the Main Theorem.” In fact, it suffices to use a simplified form of that result, which is stated in this section as the Main Lemma.

“Inductive Lemmas” and “the Main Theorem” in this section refer to the corresponding statements in [Mur05]. Because they are not used outside of this section, because their statements involve properties of maps that are not defined in this paper, and because the Main Lemma may be viewed as a particular case of the Main Theorem, the Main Theorem and Inductive Lemmas are not stated here.

Definition. Call a map Δ *semisimple* if every edge of Δ is incident to a face of Δ .

Definition. Let Δ be a semisimple S-map. Let S be the number of selected external edges of Δ . Let μ be a real number. Then the S-map Δ is said to satisfy the *condition* $\mathcal{X}(\mu)$ if

$$S \geq \|\Delta(1)\| - \mu \sum_{\Pi \in \Delta(2)} |\partial\Pi|.$$

Note that the definition of the condition $\mathcal{X}$ in [Mur05], given there only for simple S-maps, is extended here to all semisimple S-maps.

Proposition 4.1. *Let Δ be an S-map, and let μ be a real number. Suppose every maximal semisimple S-submap of Δ satisfies the condition $\mathcal{X}(\mu)$. Let S be the number of selected external edges of Δ . Then*

$$S \geq (1 - 2\mu) \sum_{\Pi \in \Delta(2)} |\partial\Pi|.$$

Proof. Consider an arbitrary maximal semisimple S-submap Δ' of Δ . Let S' be the number of selected external edges of Δ' . Let Σ' be the sum of the degrees of all the faces of Δ' . Then

$$S' = 2S' - S' \geq 2\|\Delta'(1)\| - 2\mu\Sigma' - S' \geq S' + \Sigma' - 2\mu\Sigma' - S' = (1 - 2\mu)\Sigma'.$$

Now, consider Δ itself. Remove from the underlying 2-complex of Δ all edges that are not incident to any face. All connected components of the remaining 2-complex (some, possibly, consisting of just one vertex) define

semisimple S-submaps of Δ . These S-submaps are precisely all maximal semisimple S-submaps of Δ . Therefore, every selected external edge of Δ , as well as every face of Δ , belongs to exactly one maximal semisimple S-submap of Δ . Hence, the desired inequality for Δ may be obtained by adding up the analogous inequalities for all of its maximal semisimple S-submaps. $\square$

Although it takes little effort to derive the Main Lemma formulated below from the Main Theorem, it will be even easier if a modified version of the Main Theorem with slightly weakened hypotheses is used instead. It can be observed that the Main Theorem will still hold true if “simple S-map” is replaced with “semisimple S-map” in the hypotheses. Indeed, analysis of the proofs of Inductive Lemmas shows that their statements can be modified in the following way without turning them false:

- (1) The hypotheses of Inductive Lemma 1 may be weakened by requiring the condition $\mathcal{X}(\mu)$ only from all simple disc S-submaps of Δ all of whose edges are internal in Δ , rather than from all proper simple disc S-submaps.
- (2) The hypotheses of Inductive Lemma 2 may be weakened by requiring the S-map Δ to be only semisimple, rather than simple.

The original proofs with a few obvious changes apply to the modified lemmas.

The statement obtained from the Main Theorem by replacing “simple” with “semisimple” shall be referred to as the Modified Main Theorem. The original proof of the Main Theorem applies to the Modified Main Theorem if Inductive Lemmas, used in the proof, are modified as described. (The condition $\mathcal{X}$ in the conclusions of the Modified Main Theorem and modified Inductive Lemma 2 shall be understood in the above-defined broader sense.)

Remark 4.1. The Modified Main Theorem can also be obtained as a corollary of the Main Theorem by “cutting” a given semisimple S-map into its “simple components.”

Definition. Let Δ be an S-map. Let $\lambda_1, \lambda_2 \in [0, 1]$. The S-map Δ is said to satisfy the *condition* $\mathcal{B}(\lambda_1, \lambda_2)$ if it satisfies the following three conditions:

$\mathcal{B}_0$: For each face Π of Δ , the contour of Π has at least one selected subpath and at most one maximal selected subpath (note that if all nontrivial subpaths are selected, then there is no maximal selected subpath).

$\mathcal{B}_1(\lambda_1)$: For each face Π of Δ , there is a selected subpath of $\partial\Pi$ of length at least $(1 - \lambda_1)|\partial\Pi|$.

$\mathcal{B}_2(\lambda_2)$: For each face Π of Δ , the length of every double-selected arc incident to Π is at most $\lambda_2|\partial\Pi|$.

A map Δ is said to satisfy the condition $\mathcal{B}(\lambda_1, \lambda_2)$ if there exists a selection on Δ such that Δ with this selection satisfies $\mathcal{B}(\lambda_1, \lambda_2)$.

For all admissible values of λ_1 and λ_2 , the condition $\mathcal{B}(\lambda_1, \lambda_2)$ is equivalent to the condition $\mathcal{A}(1, 1, 1; \lambda_1, \lambda_2, 0, 0)$ in [Mur05].

Main Lemma. *Let Δ be an S-map with at most 3 contours. Let $\lambda_1, \lambda_2 \in [0, 1]$. Suppose $2\lambda_1 + 13\lambda_2 < 1$ and Δ satisfies $\mathcal{B}(\lambda_1, \lambda_2)$. Let S be the*

number of selected external edge of Δ , and let $\mu = \lambda_1 + 5\lambda_2$. Then

$$S \geq (1 - 2\mu) \sum_{\Pi \in \Delta(2)} |\partial\Pi|.$$

Proof. Every maximal semisimple S-submap of Δ has at most 3 contours (see the description of maximal semisimple S-submaps given in the proof of Proposition 4.1) and satisfies the condition $\mathcal{B}(\lambda_1, \lambda_2)$. By the Modified Main Theorem, and Remarks 6.3 and 6.4 in [Mur05], all maximal semisimple S-submaps of Δ satisfy the condition $\mathcal{X}(\mu)$. The desired inequality now follows from Proposition 4.1. $\square$

5. PROPERTIES OF THE GROUP

In this section some properties of the above-defined group G are established. In particular, it is shown that G is an example that provides the negative answer to Bludov's question.

Adopt the notation of Section 2. In particular, define n and λ_1 the same way. Let $\lambda_2 = 2/n$, $\mu = \lambda_1 + 5\lambda_2$. Then

$$(5.1) \quad 2\lambda_1 + 13\lambda_2 < 1$$

and

$$(5.2) \quad 1 - 2\mu - 2\lambda_1 - \frac{2n\lambda_1^2}{1 - \lambda_1} \geq 1 - \frac{21}{n}.$$

In particular $\mu < 1/2$. It is well known that $21 \times 3 = 63$; this explains why 63.

Inequality (2.4) is equivalent to

$$(5.3) \quad m_i \geq \frac{1 - \lambda_1}{n} (nm_i + |w_i|).$$

Combining (2.4), (5.3), and (2.3), obtain

$$(5.4) \quad j \leq i \Rightarrow |w_j| \leq \frac{n\lambda_1}{1 - \lambda_1} m_i.$$

Inequality (2.1) implies that $\lambda_1 < 1/(4n) < 1/(2n + 1)$. It follows from this and from (5.4) that

$$(5.5) \quad j \leq i \Rightarrow |w_j| < \frac{m_i}{2}.$$

If w is a group word over $\mathfrak{A}$, let $[w]_G$, $[w]_{\mathcal{R}}$, or simply $[w]$, denote the element of G represented by w . Let $a_1, \dots, a_n$ be the elements of G represented by the one-letter group words $x_1, \dots, x_n$, respectively (in terms of “brackets,” $a_i = [x_i]$).

In this section a selection on a diagram Δ is called *special* if the contour of every face Π of Δ has two subpaths s and t such that:

- $\langle st \rangle = \partial\Pi$;
- s is the only maximal selected subpath of $\partial\Pi$;
- there is $m \in \mathbb{N}$ such that either $\ell(s) = x_1^m x_2^m \dots x_n^m$ or $\ell(s) = x_n^{-m} x_{n-1}^{-m} \dots x_1^{-m}$;
- $|s| > \frac{n}{2n-2} |\partial\Pi|$ (note that $|\partial\Pi| = |s| + |t|$).

On every diagram Δ over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$, there exists a unique special selection. (In fact, if Δ is a diagram over an arbitrary group presentation, and a special in the above sense selection on Δ exists, then it is unique.) A diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ together with a special selection shall be called a *special S-diagram*. Note that if s is the maximal selected subpath of the contour of a face Π of a special S-diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$, then $|s| \geq (1 - \lambda_1)|\partial\Pi|$, which is stronger than the inequality in the definition of a special selection.

If Δ is a diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$, and Π is a face of Δ , then define the *rank* of the face Π to be such j that the label of some representative of $\partial\Pi$ is $r_j^{\pm 1}$. Clearly, the rank of a face is well-defined. Let the rank of a face Π be denoted by $\text{rank}(\Pi)$. Note that if $\text{rank}(\Pi_1) \geq \text{rank}(\Pi_2)$, then $|\partial\Pi_1| \geq |\partial\Pi_2|$. The ranks of two faces in a special S-diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ are equal if and only if the lengths of the maximal selected subpaths of their contours are equal (follows from (2.2)).

Proposition 5.1. *Every weakly reduced special S-diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ satisfies the condition $\mathcal{B}(\lambda_1, \lambda_2)$.*

Proof. Let Δ be a weakly reduced special S-diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$. Clearly, Δ satisfies $\mathcal{B}_0$.

If Π is a face of Δ , s is the maximal selected subpath of $\partial\Pi$, and t is the path such that $\langle st \rangle = \partial\Pi$, then $\ell(s) = (x_1^{m_j} x_2^{m_j} \dots x_n^{m_j})^{\pm 1}$ and $\ell(t) = w_j^{\mp 1}$ where $j = \text{rank } \Pi$. Since $|w_j| \leq \lambda_1 |r_j|$ (see (2.4)), have that $|s| \geq (1 - \lambda_1)|\partial\Pi|$. Hence, Δ satisfies $\mathcal{B}_1(\lambda_1)$.

If u is a double-selected oriented arc, then $\ell(u)$ is a subword of a word of the form $x_l^m x_{l+1}^m$ or $x_{l+1}^{-m} x_l^{-m}$ (since Δ is weakly reduced). Therefore, if such a u is a subpath of the contour of a face Π , then $|u| < (2/n)|\partial\Pi| = \lambda_2 |\partial\Pi|$. Hence, Δ satisfies $\mathcal{B}_2(\lambda_2)$. $\square$

Corollary 5.1.1. *If Δ is a weakly reduced special S-diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ with at most 3 contours, and S is the number of selected external edge of Δ , then*

$$S \geq (1 - 2\mu) \sum_{\Pi \in \Delta(2)} |\partial\Pi|.$$

Proof. Follows from the proposition and the Main Lemma. $\square$

Corollary 5.1.2. *The group presentation $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ is strongly aspherical in the sense that no spherical diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ is weakly reduced.*

Proof. Every weakly reduced special spherical S-diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ has at least $(1 - 2\mu) \cdot 2$ selected external edges and therefore does not exist ($\mu < 1/2$). $\square$

Proposition 5.2. *Let Δ be a special nondegenerate S-diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$. Let $\mathfrak{B}$ be an arbitrary nonempty subset of $\mathfrak{A}$, and let $k = |\mathfrak{B}|$. Let S be the number of selected external edges of Δ whose labels are in $\mathfrak{B}$. Then*

$$S < \frac{k}{n} \sum_{\Pi \in \Delta(2)} |\partial\Pi|.$$

Proof. If Π is a face of Δ , and s is the maximal selected subpath of $\partial\Pi$, then $\ell(s) = (x_1^{m_j} x_2^{m_j} \dots x_n^{m_j})^{\pm 1}$ where $j = \text{rank } \Pi$. Therefore, the number of

selected external edges of Δ incident to Π whose labels are in $\mathfrak{B}$ is at most $(k/n)|s| < (k/n)|\partial\Pi|$. Since this is true for every face Π of Δ , the desired inequality follows. $\square$

Definition. Let $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$ be a finite group presentation. A function $f: \mathbb{N} \rightarrow \mathbb{R}$ is called an *isoperimetric function* of $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$ if for every group word w over $\mathfrak{B}$ equal to 1 modulo the relations $r = 1$, $r \in \mathcal{S}$, there exists a disc diagram Δ over $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$ with at most $f(|w|)$ faces such that w is the label of some representative of $\partial\Delta$. The minimal isoperimetric function of $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$ is called the *Dehn function* of $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$.

Proposition 5.3. *If $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$ is a finite subpresentation of $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ ($\mathfrak{B}$ is a subset of $\mathfrak{A}$, and $\mathcal{S}$ is a subset of $\mathcal{R}$), then the function $f: \mathbb{N} \rightarrow \mathbb{R}$ given by the formula*

$$f(k) = \frac{k}{1 - 2\mu}$$

is an isoperimetric function of $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$ (recall that $\mu < 1/2$).

Proof. If Δ is a weakly reduced disc diagram over $\langle \mathfrak{B} \parallel \mathcal{S} \rangle$, and $\langle q \rangle = \partial\Delta$, then, as follows from Corollary 5.1.1 of Proposition 5.1,

$$|q| \geq (1 - 2\mu) \sum_{\Pi \in \Delta(2)} |\partial\Pi| \geq (1 - 2\mu) \|\Delta(2)\|.$$

Hence, $\|\Delta(2)\| \leq f(|q|)$. $\square$

Corollary 5.3.1. *Every finite subpresentation of $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ presents a hyperbolic group.*

Proof. Use the characterization of hyperbolic groups in terms of isoperimetric functions of their finite presentations. According to Theorem 2.5, Theorem 2.12, and Corollary of the latter in [ABC⁺91], a group is hyperbolic if and only if it has a finite presentation with a linear isoperimetric function. (Note that in [ABC⁺91] all isoperimetric functions in the sense of the last definition are called “Dehn functions.”) $\square$

Let *deg-lex* be the order on the set of all group words over $\mathfrak{A}$ described as follows. Two group words are compared first by length, and second alphabetically according to the following order on the group letters: $x_1 < x_1^{-1} < x_2 < \dots < x_n < x_n^{-1}$. For example, $x_n x_n < x_1 x_1 x_1$ and $x_1 x_2 x_3 < x_1 x_3 x_2$.

Proposition 5.4. *If the order on reduced group words used in Section 2 for choosing $\{w_i\}_{i=1,\dots}$ is deg-lex, N is a positive integer, and $m_i = N|w_i| + i$ for every i for which r_i is defined, then the presentation $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ is recursive.*

Proof. If the presentation $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ is finite, then it is recursive. (It will be shown in Section 6 that it cannot be finite.) Now, assume that the presentation is infinite.

Let C be the set of all 4-tuples $(\mathcal{S}, E, u, v)$ such that $\mathcal{S}$ is a finite set of group words over $\mathfrak{A}$, E is a rational number, u and v are group words over $\mathfrak{A}$, and there exists a disc diagram over $\langle \mathfrak{A} \parallel \mathcal{S} \rangle$ with at most E edges such that the label of some representative of its contour is uv^{-1} . The set C is recursive.

Let q be a rational number such that $q \geq 1/(1 - 2\mu)$. Let D be the set of all 3-tuples $(\mathcal{S}, u, v)$ such that

$$\left(\mathcal{S}, \frac{1+qL}{2}(|u| + |v|), u, v\right) \in C$$

if L is the maximum of the lengths of the elements of $\mathcal{S}$. The set D is recursive. This set describes consequences of a given finite set of relations as explained below.

Let f be the function $\mathbb{N} \rightarrow \mathbb{R}, k \mapsto qk$. Consider an arbitrary finite $\mathcal{S}$ such that f is an isoperimetric function of $\langle \mathfrak{A} \parallel \mathcal{S} \rangle$. Let u and v be arbitrary group words over $\mathfrak{A}$. Then the relation $u = v$ is a consequence of the relations of $\langle \mathfrak{A} \parallel \mathcal{S} \rangle$ if and only if $(\mathcal{S}, u, v) \in D$.

Consider the following algorithm:

Input: a positive integer j .
 Step 1: Produce the set $\mathcal{S}$ of outputs of this algorithm on the inputs $1, \dots, j-1$. (Do not stop if the algorithm does not stop on at least one of those inputs; if $j=1$, then $\mathcal{S} = \emptyset$.) Let L be the maximum of the lengths of the elements of $\mathcal{S}$ if $\mathcal{S}$ is nonempty, and 0 otherwise.
 Step 2: Find the least (with respect to deg-lex) group word w over $\mathfrak{A}$ such that for every regular group word u of length at most $(n+1)|w| + n^4L$, $(\mathcal{S}, u, w) \notin D$. (Do not stop if there is no such w .) Let $m = N|w| + j$.
 Output: $x_1^m x_2^m \dots x_n^m w^{-1}$.

It shall be shown by induction that on every input $j \in \mathbb{N}$, the algorithm stops and gives r_j as the output. It will follow that $\mathcal{R}$ is recursive because the length of r_j is an increasing function of j .

Take an arbitrary $k \in \mathbb{N}$. If $k > 1$, assume that it is already proved that for every positive integer $j < k$, the algorithm gives r_j as the output on the input j . Consider the work of the algorithm on the input k .

By the inductive assumption, the set $\mathcal{S}$ produced in Step 1 coincides with $\mathcal{R}_{k-1}$. In particular, f is an isoperimetric function of $\langle \mathfrak{A} \parallel \mathcal{S} \rangle$. Note that $L = |r_{k-1}|$ if $k > 1$, and $L = 0$ if $k = 1$.

By the choice of w_k , this word is not equal to any regular group word modulo the relations of $\langle \mathfrak{A} \parallel \mathcal{S} \rangle$. Therefore, Step 2 of the algorithm is carried out in finite time, and the obtained group word w is not greater than w_k with respect to the deg-lex order.

To complete the inductive step and the proof of this proposition, it suffices to show that the group word w obtained in Step 2 is w_k . Suppose w is distinct from w_k .

Since $w < w_k$ relative to deg-lex, it follows from the choice of w_k that w equals some regular group word modulo the relations of $\langle \mathfrak{A} \parallel \mathcal{S} \rangle$. Let u be such a regular group word. By the choice of w ,

$$|u| > (n+1)|w| + n^4L.$$

Let Δ be a weakly reduced special disc S-diagram over $\langle \mathfrak{A} \parallel \mathcal{S} \rangle$ such that the label of some representative of $\partial\Delta$ is uw^{-1} . Let b and p be the paths such that $\ell(b) = u$, $\ell(p) = w^{-1}$, and $\langle bp \rangle = \partial\Delta$. Let $b_1, b_2, \dots, b_n$ be the subpaths of b such that $b = b_1 b_2 \dots b_n$, and for every $i = 1, 2, \dots, n$, the label of b_i is a power of x_i . For every $i = 1, 2, \dots, n$, let B_i be the set of all edges that lie on b_i . Let Σ be the sum of the degrees of all the faces of Δ . Let T be the number of selected external edges of Δ that lie on b .

Since $\ell(b)$ is regular, no two oriented edges of b are inverse to each other. Hence,

$$|b| \leq \Sigma + |p|.$$

Let Σ_1 be the sum of the degrees of all faces of Δ that are incident with edges from at least 3 distinct sets from among $B_1, \dots, B_n$. If $1 \leq i_1 < i_2 < i_3 \leq n$, then there is at most one face that is incident to edges from all three sets $B_{i_1}, B_{i_2}, B_{i_3}$. Therefore, $\Sigma_1 \leq n^3 L$ (the degree of every face of Δ is at most L). Let Σ_2 be the sum of the degrees of all faces of Δ that are incident with edges from no more than 2 distinct sets from among $B_1, \dots, B_n$.

One one hand,

$$T \leq \Sigma_1 + \frac{2}{n} \Sigma_2 \leq n^3 L + \frac{2}{n} \Sigma.$$

On the other hand, by Proposition 5.1 and the Main Lemma,

$$T + |p| \geq (1 - 2\mu)\Sigma.$$

Therefore,

$$\begin{aligned} (1 - 2\mu)\Sigma &\leq n^3 L + \frac{2}{n} \Sigma + |p|, \\ \left(1 - 2\mu - \frac{2}{n}\right)\Sigma &\leq |p| + n^3 L, \\ \frac{n - 23}{n} \Sigma &\leq |p| + n^3 L, \\ \Sigma &\leq n|p| + n^4 L, \\ |b| &\leq (n + 1)|p| + n^4 L. \end{aligned}$$

In other terms, $|u| \leq (n + 1)|w| + n^4 L$. This gives a contradiction.

Thus, $w = w_k$, and the inductive step is done. $\square$

Everything is now ready to start proving properties of G .

Property 1. *For every $(n - 21)$ -element subset I of the set $\{1, \dots, n\}$, the system $\{a_i\}_{i \in I}$ freely generates a free subgroup F of G (of rank $n - 21$). Moreover, every two elements in such a free subgroup F are conjugate in G only if they are conjugate in F .*

Proof. Suppose that Property 1 does not hold. Take a set $I \subset \{1, \dots, n\}$ of $n - 21$ elements that provides a counterexample. Let $\mathfrak{B} = \{x_i \mid i \in I\}$.

If the system $\{a_i\}_{i \in I}$ does not freely generate a free subgroup of G , then there is a weakly reduced disc diagram Δ over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ such that the label of some (therefore every) representative of its contour is a cyclically reduced nonempty group word over the alphabet $\mathfrak{B}$. Clearly, such a diagram Δ is nondegenerate (has a face).

If the system $\{a_i\}_{i \in I}$ does freely generate a free subgroup F of G , but there are two elements of F that are conjugate in G but not in F , then

let v_1 and v_2 be two cyclically reduced group words over $\mathfrak{B}$ that represent two such elements. The group words v_1 and v_2 are not cyclic shifts of each other, and neither of them represents the identity of G . Since $[v_1]$ and $[v_2]$ are nontrivial but are conjugate in G , there exists a weakly reduced annular diagram Δ over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ such that the label of some representative of one of its contours is v_1 , and the label of some representative of its other contour is v_2^{-1} (see Lemma V.5.2 in [LS01]). Clearly, such Δ is nondegenerate.

Thus, to obtain a contradiction and complete the proof, it is enough to show that there is no weakly reduced nondegenerate disc nor annular diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ with fewer than $n - 20$ distinct basic letters on its contour(s).

Let Δ be an arbitrary special weakly reduced nondegenerate disc or annular S-diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$. (Recall that a special selection exists on every diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$.) Suppose that Δ has fewer than $n - 20$ distinct basic letters of $\mathfrak{A}$ on its contour(s). Let S be the total number of selected external edges of Δ . On one hand, by Proposition 5.2,

$$S < \frac{n - 21}{n} \sum_{\Pi \in \Delta(2)} |\partial \Pi|.$$

On the other hand, by Corollary 5.1.1 of Proposition 5.1,

$$S \geq (1 - 2\mu) \sum_{\Pi \in \Delta(2)} |\partial \Pi|.$$

Since

$$1 - 2\mu > \frac{n - 21}{n},$$

one has a contradiction. $\square$

The proof of the next property is not so straightforward.

Property 2. *For every element g of G , there exist unique $k_1, \dots, k_n \in \mathbb{Z}$ such that $g = a_1^{k_1} \dots a_n^{k_n}$.*

The uniqueness is the “hard part” of Property 2. It shall be proved by contradiction. Main steps of the proof are stated below as Lemmas 5.1–5.16. These lemmas share some common assumptions about an S-diagram Δ .

Assume that Δ is a special disc S-diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ whose contour has a representative of the form $p_1 p_2$ such that $\ell(p_1)$ and $\ell(p_2^{-1})$ are distinct regular group words, and $\ell(p_1 p_2)$ is cyclically reduced. Suppose, moreover, that Δ is such an S-diagram with the minimal possible number of faces. In particular, if Δ' is a nontrivial disc diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$, p is a representative of $\partial \Delta'$, and $\ell(p)$ is regular or counter-regular, then Δ' has no fewer faces than Δ .

Lemma 5.1. *The diagram Δ is weakly reduced.*

Proof. This easily follows from the minimality of the number of faces of Δ . $\square$

Lemma 5.2. *The diagram Δ is simple.*

Proof. Use the minimality of Δ again. Some maximal simple disc S -subdiagram of Δ satisfies all the assumptions made about Δ (this follows, for example, from Proposition 3.1 of [Mur05]), and hence it must be the whole of Δ . $\square$

Lemma 5.3. *If Δ' is a disc S -subdiagram of Δ , and S is the number of selected external edge of Δ' (external in Δ'), then*

$$S \geq (1 - 2\mu) \sum_{\Pi \in \Delta'(2)} |\partial\Pi|.$$

Proof. Follows directly from Lemma 5.1 and Corollary 5.1.1 of Proposition 5.1. $\square$

Lemma 5.4. *If Δ' is a disc subdiagram of Δ , p_1 and p_2 are paths, $\langle p_1 p_2 \rangle = \partial\Delta'$, and the reduced forms of $\ell(p_1)$ and $\ell(p_2^{-1})$ are regular, then either Δ' is degenerate, or it coincides with Δ .*

Proof. Suppose Δ' , p_1 , and p_2 are such as in the hypotheses of the lemma.

Case 1: $\ell(p_1)$ and $\ell(p_2^{-1})$ are freely equal. Then the label of every representative of $\partial\Delta'$ is freely trivial and therefore Δ' is degenerate. Indeed, if Δ' was nondegenerate, there would exist a disc diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ with the same label of the contour as Δ but with fewer faces (faces of Δ' , possibly together with some of the others, could be “eliminated” from Δ), which would contradict the minimality of Δ .

Case 2: $\ell(p_1)$ and $\ell(p_2^{-1})$ are not freely equal. Then take Δ' and repeatedly *fold* labelled external edges and “cut off,” whenever necessary, “branches” with freely trivial contour labels until one obtains a disc diagram Δ'' such that the labels of the representatives of $\partial\Delta''$ are cyclically reduced. Clearly, Δ'' is a nondegenerate diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$, the label of every representative of $\partial\Delta''$ is a cyclically reduced form of $\ell(p_1 p_2)$, and $\|\Delta''(2)\| \leq \|\Delta'(2)\|$. By the minimality of Δ , such a diagram Δ'' cannot have fewer faces than Δ . Therefore, $\Delta'(2) = \Delta(2)$ and, since $\partial\Delta$ is reduced, Δ' coincides with Δ . $\square$

Lemma 5.5. *The diagram Δ has more than 1 face.*


Proof. This easily follows from the form of elements of $\mathcal{R}$. $\square$

Lemma 5.6. *There are at least $n - 20$ distinct basic letters on $\partial\Delta$.*

Proof. Follows from Property 1. $\square$

Let b_1 be the maximal subpath of $\partial\Delta$ whose label is regular but not counter-regular (is not a letter power). Let b_2 be the maximal subpath of $\partial\Delta$ whose label is counter-regular but not regular. Since, according to Lemma 5.6, there are at least 2 distinct basic letters on $\partial\Delta$, the paths b_1 and b_2 are well-defined. (If, say, the label of some representative of $\partial\Delta$ is $x_1 x_2 x_3 \dots x_{42} x_{43} x_{42} x_2 x_1$, then $\ell(b_1) = x_1^2 x_2 \dots x_{43}$, and $\ell(b_2) = x_{43} x_{42} x_2 x_1^2$.) The label of each of the paths b_1 and b_2 has at least 2 distinct basic letters. Let l_1 be the initial subpath of b_1 that is a terminal subpath of b_2 , and let l_2 be the initial subpath of b_2 that is a terminal subpath of b_1 . Both l_1 and l_2 are nontrivial (see Fig. 1). The labels of l_1 and l_2 are letter powers.

Lemma 5.7. *Every selected external arc of Δ lies on at least one of the paths b_1 or b_2 .*

FIGURE 1. The paths b_1 , b_2 , l_1 , and l_2 in Δ .

Proof. Follows from the fact that the label of every selected oriented arc is either regular or counter-regular. $\square$

The following observation is obvious but deserves mentioning because it is used implicitly in the proofs of Lemmas 5.8, 5.10, and 5.13 several times.

Suppose p_1 , p_2 , and q are paths in a diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$, and q is nontrivial. Then

- if the products p_1q and $q^{-1}p_2$ are defined and their labels are regular, then $\ell(q)$ is a letter power, and the reduced form of $\ell(p_1p_2)$ is regular;
- if the products p_1q and qp_2 are defined and their labels are regular, then $\ell(p_1qp_2)$ is regular.

Since Lemma 5.7 and many of the remaining lemmas deal with selected external arcs or selected external oriented arcs, it is advisable to review these concepts. An oriented arc of a graph (or a 2-complex) is a simple path all of whose vertices, except, possibly, the end-vertices, have degree two in the given graph. A (non-oriented) arc is a pair of mutually inverse oriented arcs. An arc is incident to a face Π if (and only if) at least one of the associated oriented arcs is a subpath of $\partial\Pi$. Every external arc of a (semi-)simple map is incident to exactly one face. An oriented arc is selected if (and only if) it is selected as a path (in particular, it must be a subpath of the contour of some face). An external arc is selected if (and only if) it is incident to some face Π , and the associated oriented arc that is a subpath of $\partial\Pi$ is selected. Distinct maximal selected external arcs of Δ never overlap.

Lemma 5.8. *In the diagram Δ , every face is incident with at most 1 maximal selected external arc.*

Proof. Consider an arbitrary face Π of Δ . Suppose Π is incident with at least 2 distinct maximal selected external arcs.

Let u_1 and u_2 be distinct maximal selected external oriented arcs of Δ that are subpaths of $\partial\Pi$. (They do not overlap.) Let s be the maximal selected subpath of $\partial\Pi$. Without loss of generality, assume that u_1 precedes

u_2 as a subpath of s . Let v be such a path that u_1vu_2 is a subpath of s . Since Δ is simple (see Lemma 5.2), and u_1 and u_2 are maximal, it follows that v is nontrivial, and the first and last oriented edges of v are internal in Δ .

Suppose v has a simple cyclic subpath p . Then $\langle p \rangle$ is the contour of a proper simple disc subdiagram of Δ . Since $\ell(p)$ is regular or counter-regular, this contradicts the minimality of Δ . Hence, the path v is simple and not cyclic (equivalently, every vertex occurs in v at most once).

Let v_1 be the minimal subpath of $\partial\Delta$ such that u_1^{-1} and u_2^{-1} are respectively its initial and terminal subpaths. Let v_2 be the path such that $\langle v_1v_2 \rangle = \partial\Delta$. The paths v_1v^{-1} and v_2v are cyclically reduced and represent the contours of two disc subdiagrams of Δ . Let Δ'_1 and Δ'_2 be the subdiagrams with the contours $\langle v_1v^{-1} \rangle$ and $\langle v_2v \rangle$, respectively. Note that $\Pi \in \Delta'_1(2)$, and $\Delta(2)$ is the disjoint union of $\Delta'_1(2)$ and $\Delta'_2(2)$. The diagram Δ'_1 is simple. The diagram Δ'_2 is nondegenerate. Both Δ'_1 and Δ'_2 are proper subdiagrams of Δ .

The labels of u_1 and u_2 are both regular or both counter-regular. Therefore, each of the paths u_1^{-1} and u_2^{-1} is a subpath of one of the paths b_1 or b_2 (see Lemma 5.7). Consider the following 4 cases (see Fig. 2; shaded subdiagrams are the “sources of contradiction”):

Case 1: u_1^{-1} and u_2^{-1} are subpaths of b_1 . Then either v_1 or $u_2^{-1}v_2u_1^{-1}$ is a subpath of b_1 as well. Suppose v_1 is a subpath of b_1 . Then $\ell(v_1)$ is regular, $\ell(u_1)$ and $\ell(u_2)$ are letter powers, and $\ell(u_1vu_2)$ is regular. Therefore, $\langle v_1v^{-1} \rangle$ cannot be the contour of a proper nondegenerate disc subdiagram (see Lemma 5.4). This gives a contradiction. Suppose $u_2^{-1}v_2u_1^{-1}$ is a subpath of b_1 . Then $\ell(u_2^{-1}v_2u_1^{-1})$ is regular, and $\ell(u_1vu_2)$ is counter-regular. Therefore, $\langle v_2v \rangle$ cannot be the contour of a proper nondegenerate disc subdiagram. This gives a contradiction.

Case 2: u_1^{-1} and u_2^{-1} are subpaths of b_2 . When proving impossibility of Case 1, it was essentially shown that no face of Δ can be incident with 2 distinct maximal selected external arcs lying on b_1 . Since all the assumptions made about Δ hold for its mirror copy as well, the same statement appropriately reformulated must hold for the mirror copy of Δ . Namely, no face of the mirror copy of Δ can be incident with 2 distinct maximal selected external arcs lying on b_2^{-1} (b_2^{-1} plays the same role for the mirror copy of Δ as b_1 does for Δ). This means that Case 2 is impossible.

Case 3: u_1^{-1} is a subpath of b_1 , and u_2^{-1} is a subpath of b_2 . Pick a subpath p_1 of b_1 and a subpath p_2 of b_2 such that u_1^{-1} is an initial subpath of p_1 , u_2^{-1} is a terminal subpath of p_2 , and $p_1p_2 = v_1$. Then $\ell(p_1)$ is regular, $\ell(p_2)$ is counter-regular, and $\ell(u_2^{-1}v^{-1}u_1^{-1})$ is regular or counter-regular. If $\ell(u_2^{-1}v^{-1}u_1^{-1})$ is regular, then $\ell(v^{-1}p_1)$ is regular. If $\ell(u_2^{-1}v^{-1}u_1^{-1})$ is counter-regular, then $\ell(p_2v^{-1})$ is counter-regular. In either case $\langle p_2v^{-1}p_1 \rangle = \langle v_1v^{-1} \rangle$ cannot be the contour of a proper nondegenerate disc subdiagram. This gives a contradiction.

Case 4: u_1^{-1} is a subpath of b_2 , and u_2^{-1} is a subpath of b_1 . Pick a subpath p_1 of b_2 and a subpath p_2 of b_1 such that u_1^{-1} is an initial subpath of p_1 , u_2^{-1} is a terminal subpath of p_2 , and $p_1p_2 = v_1$. Then $\ell(p_1)$ is counter-regular, $\ell(p_2)$ is regular, and $\ell(u_2^{-1}v^{-1}u_1^{-1})$ is regular or counter-regular. If $\ell(u_2^{-1}v^{-1}u_1^{-1})$




FIGURE 2. Cases 1–4, Lemma 5.8.

is regular, then $\ell(p_2 v^{-1})$ is regular. If $\ell(u_2^{-1} v^{-1} u_1^{-1})$ is counter-regular, then $\ell(v^{-1} p_1)$ is counter-regular. In either case $\langle p_2 v^{-1} p_1 \rangle = \langle v_1 v^{-1} \rangle$ cannot be the contour of a proper nondegenerate disc subdiagram. This gives a contradiction.

(Case 4 is symmetric with Case 3, but this symmetry is not simply the mirror symmetry as between Cases 1 and 2. Say, if Case 4 was indeed the case for Δ , then the mirror copy of Δ would be just another example for the same Case 4, not Case 3.)

A contradiction is obtained in each of the 4 cases, and no other cases exist. $\square$

Lemma 5.9. *There are a face Π and a path s' in Δ such that:*

- s' is a selected subpath of $\partial\Pi$;
- s' is a maximal selected external oriented arcs of Δ ;
- the $n - 40$ basic letters $x_{21}, x_{22}, \dots, x_{n-20}$ all occur in $\ell(s')$.

Proof. For every face Π of Δ , let $S(\Pi)$ denote the number of selected external edges of Δ incident to Π . By Lemma 5.3,

$$\sum_{\Pi \in \Delta(2)} S(\Pi) \geq (1 - 2\mu) \sum_{\Pi \in \Delta(2)} |\partial\Pi|.$$

Hence, there exists a face Π of Δ such that

$$S(\Pi) \geq (1 - 2\mu)|\partial\Pi| > \frac{n - 21}{n}|\partial\Pi|$$

(recall that $1 - 2\mu > 1 - 21/n$).

Let Π be such a face as above. Let s be the maximal selected subpath of $\partial\Pi$. Let s' be the (only) maximal selected external oriented arc of Δ that is a subpath of s (see Lemma 5.8). Then

$$|s'| = S(\Pi) > \frac{n - 21}{n}|\partial\Pi| > \frac{n - 21}{n}|s|.$$

Let $j = \text{rank}(\Pi)$. Then either $\ell(s) = x_1^{m_j} x_2^{m_j} \dots x_n^{m_j}$ or $\ell(s) = x_n^{-m_j} x_{n-1}^{-m_j} \dots x_1^{-m_j}$. Therefore, $\ell(s')$ has at least $n - 20$ distinct basic letters, and all of the basic letters $x_{21}, x_{22}, \dots, x_{n-20}$ occur in it. $\square$

Lemmas 5.1–5.9, as well as 5.10–5.16, assert some properties of Δ . Observe that none of these properties in fact can distinguish between Δ and its mirror copy, i.e., each of these properties holds for Δ if and only if it holds for the mirror copy of Δ . Thus, at this point analogues of Lemmas 5.1–5.9 for the mirror copy of Δ shall be assumed proved. Moreover, the initial assumptions about the S-diagram Δ are also true about its mirror copy.

Even though the statement of Lemma 5.7 is about the S-diagram Δ and the paths b_1 and b_2 , it still may be viewed as an assertion of a property of Δ because, according to the way they are chosen, b_1 and b_2 are uniquely determined for the given Δ . Therefore, the analog of Lemma 5.7 for the mirror copy of Δ states:


Every selected external arc of the mirror copy of Δ lies on at least one of the paths b_2^{-1} or b_1^{-1} .

In the proofs of Lemmas 5.10–5.13, it is convenient in some cases to pass to the mirror copy of Δ to reduce the number of cases to consider.

Lemma 5.10. *Let Π_1 be a face of Δ incident to a selected external edge of Δ , and let Π_2 be another face of Δ . Let s_1 and s_2 be the maximal selected subpaths of $\partial\Pi_1$ and $\partial\Pi_2$, respectively. Let s'_1 be the maximal selected external oriented arc of Δ that is a subpath of s_1 . Let s'_{1-} and s'_{1+} be the paths such that $s_1 = s'_{1-} s'_{1+}$. Let q_1 be the path such that $\langle s'^{-1}_1 q_1 \rangle = \partial\Delta$. Suppose there are at least 2 distinct basic letters in $\ell(s'_1)$. Suppose the paths s_2 and $(s'_{1-} q_1 s'_{1+})^{-1}$ have a common oriented edge. Then they have exactly one maximal common nontrivial subpath.*

Proof. Observe that to prove that there is exactly one maximal common nontrivial subpath of s_2 and $(s'_{1-} q_1 s'_{1+})^{-1}$, it suffices to prove the same for s_2^{-1} and $s'_{1-} q_1 s'_{1+}$. This justifies passing to the mirror copy of Δ .

Since the paths s_2 and $(s'_{1-} q_1 s'_{1+})^{-1}$ have a common oriented edge, they have at least one maximal common nontrivial subpath.

FIGURE 3. The face Π_1 in Δ , Lemma 5.10.

The label of s'_1 is regular or counter-regular. Therefore, s'_1 is a subpath of b_1^{-1} or b_2^{-1} . If it is a subpath of b_2^{-1} but not of b_1^{-1} , pass from Δ , b_1 , b_2 , l_1 , l_2 , s_1 , s_2 , s'_{1-} , s'_{1+} , s'_1 , q_1 to the mirror copy of Δ , b_2^{-1} , b_1^{-1} , l_1^{-1} , l_2^{-1} , s_1^{-1} , s_2^{-1} , s'_{1+} , s'_{1-} , s_1^{-1} , q_1^{-1} , respectively. Hence, it may and shall be assumed that s'_1 is a subpath of b_1^{-1} (see Fig. 3).

Let t_1 be the path such that $\langle s_1 t_1 \rangle = \partial \Pi_1$. Let y_- and y_+ be the paths such that $b_1 = y_- s'^{-1}_{1-} y_+$. Let b'_2 be the subpath of b_2 such that $\langle b_1 b'_2 \rangle = \partial \Delta$. Then $b_2 = l_2 b'_2 l_1$ and $q_1 = y_+ b'_2 y_-$.

Since $\ell(b_1^{-1})$ is counter-regular, and $\ell(s'_1)$ is not a letter power (it has at least 2 distinct basic letters), $\ell(s_1)$ is counter-regular as well. Hence, $\ell(s'^{-1}_{1+} s'^{-1}_{1-} y_+)$ and $\ell(y_- s'^{-1}_{1-} s'^{-1}_{1-})$ are regular.

Let $j_1 = \text{rank}(\Pi_1)$. Then $\ell(s_1) = x_n^{-m_{j_1}} x_{n-1}^{-m_{j_1}} \dots x_1^{-m_{j_1}}$ and $\ell(t_1) = w_{j_1}$.

Since $\ell(s'_1)$ has at least 2 distinct basic letters, $\ell(y_- s'^{-1}_{1+})$ and $\ell(s'^{-1}_{1-} y_+)$ have disjoint sets of basic letters. This also implies that l_1 is a proper initial subpath of $y_- s'^{-1}_{1-}$, and l_2 is a proper terminal subpath of $s'^{-1}_{1-} y_+$.

Suppose s_2 and $(s'_{1-} q_1 s'_{1+})^{-1}$ have at least two distinct maximal common nontrivial subpaths. Then let u_1 and u_2 be two paths such that:

- u_1 and u_2 are nontrivial subpaths of distinct maximal common subpaths of s_2 and $(s'_{1-} q_1 s'_{1+})^{-1}$,
- u_1 precedes u_2 as a subpath of s_2 , and
- each of the paths u_1 and u_2 is a maximal common subpath of s_2 and one of the paths y_-^{-1} , y_+^{-1} , s'^{-1}_{1-} , s'^{-1}_{1+} , or b'^{-1}_2 .

Clearly, such u_1 and u_2 exist and are non-overlapping oriented arcs of Δ . Let v be the path such that $u_1 v u_2$ is a subpath of s_2 . Since u_1 and u_2 are subpaths of distinct maximal common subpaths of s_2 and $(s'_{1-} q_1 s'_{1+})^{-1}$, the path $u_1 v u_2$ is not a subpath of $(s'_{1-} q_1 s'_{1+})^{-1}$.

Consider the following 16 cases (see Fig. 4–8):

(Notice that in a certain sense Cases 6, 8, 10, 12, 14, and 16 are “symmetric” with Cases 5, 7, 9, 11, 13, and 15, respectively.)

Case 1: each of the paths u_1 and u_2 is a subpath of q_1^{-1} . Then u_1 and




FIGURE 4. Cases 2, 3, 4, Lemma 5.10.

u_2 are external selected oriented arcs of Δ . Since there is only one maximal selected external arc of Δ incident to Π_2 (see Lemma 5.8), the path u_1vu_2 is a subpath of q_1^{-1} . This gives a contradiction.

Case 2: both u_1 and u_2 are subpaths of one of the paths s'_{1-}^{-1} or s'_{1+}^{-1} . Let z be one of the paths s'_{1-} or s'_{1+} such that both u_1 and u_2 are subpaths of z^{-1} . Then $\ell(z)$ is counter-regular. Suppose u_2 precedes u_1 as a subpath of z^{-1} . Let p be the path such that $u_1^{-1}pu_2^{-1}$ is a subpath of z . Then $\ell(u_1^{-1}pu_2^{-1})$ is counter-regular. Therefore, $\ell(u_2^{-1}v^{-1}u_1^{-1})$ is regular. The cycle $\langle pu_2^{-1}v^{-1}u_1^{-1} \rangle$ is the contour of a disc subdiagram of Δ not containing the face Π_1 but containing the face Π_2 . This contradicts the minimality of Δ . Therefore, u_1 precedes u_2 as a subpath of z^{-1} . Let p be the path such that $u_2^{-1}pu_1^{-1}$ is a subpath of z . Then $\ell(u_2^{-1}pu_1^{-1})$ is counter-regular. Therefore, $\ell(u_1vu_2)$ is regular. The cycle $\langle pv \rangle$ is the contour of a disc subdiagram of Δ not containing the faces Π_1 and Π_2 . By Lemma 5.4 (by the minimality of Δ), this subdiagram is degenerate. Since z and v are reduced, $v = p^{-1}$. Therefore, u_1vu_2 is a subpath of z^{-1} . This gives a contradiction.

Case 3: u_1 is a subpath of s'_{1-}^{-1} , and u_2 is a subpath of s'_{1+}^{-1} . Let p be the path such that $u_1^{-1}p^{-1}u_2^{-1}$ is a subpath of s_1 . Then $\ell(u_2pu_1)$ is regular. Therefore, $\ell(u_1vu_2)$ is counter-regular. The cycle $\langle pu_1vu_2 \rangle$ is the contour of a disc subdiagram of Δ containing the face Π_1 but not containing the face Π_2 . This contradicts the minimality of Δ (see Lemma 5.4).

Case 4: u_1 is a subpath of s'_{1+}^{-1} , and u_2 is a subpath of s'_{1-}^{-1} . Since the labels of s'_{1-} and s'_{1+} have disjoint sets of basic letters, $\ell(s_2)$ is regular. Let p be the path such that $u_2^{-1}p^{-1}u_1^{-1}$ is a subpath of s_1 . Then $\ell(u_1vu_2)$




FIGURE 5. Special subcases of Case 4, Lemma 5.10.

and $\ell(u_1pu_2)$ are regular. If v is trivial, then the cyclic path $pv^{-1} = p$ is cyclically reduced because $\ell(p)$ is regular. If v is nontrivial, then the cyclic path pv^{-1} is cyclically reduced by the maximality of u_1 and u_2 , and because the paths v and p are reduced. Suppose the path pv^{-1} is not simple. Then some subpath of some cyclic shift of vp^{-1} is a representative of the contour of a proper simple disc subdiagram of Δ . Let Δ' be such a subdiagram. At least one of the following three subcases takes place:

- (a) there is a subpath of v or p^{-1} which represents $\partial\Delta'$, or
- (b) there are paths v' and p' such that $\langle v'p'^{-1} \rangle = \partial\Delta'$, and the paths v' and p' are either terminal subpaths of v and p , respectively, or initial subpaths of v and p , respectively, or
- (c) there are an initial subpath p'_- and a terminal subpath p'_+ of the path p such that $\langle vp'^{-1}_+p'^{-1}_- \rangle = \partial\Delta'$.

In subcase (c) note that $\ell(p'_-p'_+)$ is regular. Lemma 5.4 easily yields a contradiction in each subcase. Hence, the cyclic path pv^{-1} is simple. Therefore, it represents the contour of a simple disc subdiagram of Δ containing the faces Π_1 and Π_2 . By Lemma 5.4, $\langle pv^{-1} \rangle = \partial\Delta$. Therefore, $v = q_1^{-1}$, and u_1vu_2 is a subpath of $(s'_{1-}q_1s'_{1+})^{-1}$. This gives a contradiction.

Case 5: u_1 is a subpath of s'^{-1}_{1-} , and u_2 is a subpath of y^{-1}_- . Let p be the path such that $u_2^{-1}pu_1$ is a subpath of $y_-s'^{-1}_1s'^{-1}_{1-}$. Then $\ell(u_2^{-1}pu_1)$ is regular. Therefore, $\ell(u_1vu_2)$ is counter-regular. The cycle $\langle pu_1v \rangle$ is the contour of a disc subdiagram of Δ containing the face Π_1 but not containing the face Π_2 . This contradicts the minimality of Δ (see Lemma 5.4).

Case 6: u_1 is a subpath of y^{-1}_+ , and u_2 is a subpath of s'^{-1}_{1+} . Let p be the path such that $u_2pu_1^{-1}$ is a subpath of $s'^{-1}_{1+}s'^{-1}_{1-}y_+$. Then $\ell(u_2pu_1^{-1})$ is regular. Therefore, $\ell(u_1vu_2)$ is counter-regular. The cycle $\langle pvu_2 \rangle$ is the contour of a




FIGURE 6. Cases 5–8, Lemma 5.10.

disc subdiagram of Δ containing the face Π_1 but not containing the face Π_2 . This contradicts the minimality of Δ .

Case 7: u_1 is a subpath of y_-^{-1} , and u_2 is a subpath of s'_{1-}^{-1} . Let p be the path such that $u_1^{-1}pu_2$ is a subpath of $y_-s'_{1-}^{-1}s'_{1-}^{-1}$. The cyclic path $pv^{-1}u_1^{-1}$ is cyclically reduced. Suppose it is not simple. Then at least one of the following two subcases takes place:

- (a) there is a subpath of v or p^{-1} which represents the contour of a proper simple disc subdiagram of Δ , or
- (b) there are paths v' and p' such that $\langle v'p'^{-1} \rangle$ is the contour of a proper simple disc subdiagram of Δ , and v' and p' are terminal subpaths of v and p , respectively.

Lemma 5.4 yields a contradiction in both subcases. Hence, the cyclic path $pv^{-1}u_1^{-1}$ is simple. Therefore, it represents the contour of a disc subdiagram of Δ containing the faces Π_1 and Π_2 . The label of $u_1^{-1}pu_2$ is regular. Therefore, $\ell(u_2^{-1}v^{-1}u_1^{-1})$ is counter-regular. By Lemma 5.4, $\langle pv^{-1}u_1^{-1} \rangle = \partial\Delta$. Therefore, u_1vu_2 is a subpath of $(s'_{1-}q_1)^{-1}$. This gives a contradiction.

Case 8: u_1 is a subpath of s'_{1+}^{-1} , and u_2 is a subpath of y_+^{-1} . Arguing as in Case 7, one obtains that u_1vu_2 is a subpath of $(q_1s'_{1+})^{-1}$. This gives a contradiction.

Case 9: u_1 is a subpath of s'_{1-}^{-1} , and u_2 is a subpath of y_+^{-1} . Let p_1 and p_2 be the paths such that $u_1^{-1}p_1$ is a terminal subpath of s'_{1-} , and $p_2u_2^{-1}$ is an initial subpath of y_+ . Then $\ell(u_1^{-1}p_1)$ is counter-regular, $\ell(p_2u_2^{-1})$ is regular, and $\ell(u_2^{-1}v^{-1}u_1^{-1})$ is regular or counter-regular. If $\ell(u_2^{-1}v^{-1}u_1^{-1})$ is regular, then $\ell(p_2u_2^{-1}v^{-1}u_1^{-1})$ is regular. If $\ell(u_2^{-1}v^{-1}u_1^{-1})$ is counter-regular,




FIGURE 7. Cases 9–12, Lemma 5.10.

then $\ell(u_2^{-1}v^{-1}u_1^{-1}p_1)$ is counter-regular. The cycle $\langle p_1p_2u_2^{-1}v^{-1}u_1^{-1} \rangle$ is the contour of a disc subdiagram of Δ containing the face Π_2 but not containing the face Π_1 . This contradicts the minimality of Δ .

Case 10: u_1 is a subpath of y_-^{-1} , and u_2 is a subpath of s'_{1+}^{-1} . Contradiction is obtained as in Case 9.

Case 11: u_1 is a subpath of y_+^{-1} , and u_2 is a subpath of s'_{1-}^{-1} . Let p_1 and p_2 be the paths such that $u_2^{-1}p_1$ is a terminal subpath of s'_{1-} , and $p_2u_1^{-1}$ is an initial subpath of y_+ . Then $\ell(u_2^{-1}p_1)$ is counter-regular, $\ell(p_2u_1^{-1})$ is regular, and $\ell(u_1vu_2)$ is regular or counter-regular. If $\ell(u_1vu_2)$ is regular, then the reduced form of $\ell(p_2v)$ is regular. If $\ell(u_1vu_2)$ is counter-regular, then the reduced form of $\ell(vp_1)$ is counter-regular. The cycle $\langle p_1p_2v \rangle$ is the contour of a disc subdiagram of Δ not containing the faces Π_1 and Π_2 . By Lemma 5.4, $\langle p_1p_2v \rangle$ is the contour of a degenerate disc subdiagram. Since $s'_{1-}y_+$ and v are reduced, $v = (p_1p_2)^{-1}$. Therefore, u_1vu_2 is a subpath of $(s'_{1-}y_+)^{-1}$. This gives a contradiction.

Case 12: u_1 is a subpath of s'_{1+}^{-1} , and u_2 is a subpath of y_-^{-1} . Arguing as in Case 11, one obtains that u_1vu_2 is a subpath of $(y_-s'_{1+})^{-1}$. This gives a contradiction.

Case 13: u_1 is a subpath of s'_{1-}^{-1} , and u_2 is a subpath of $b_2'^{-1}$. Let p_1 and p_2 be the paths such that p_1u_1 is an initial subpath of $y_-s'_{1-}^{-1}s'_{1-}^{-1}$, and $u_2^{-1}p_2$ is a terminal subpath of b_2' . Then $\ell(p_1u_1)$ is regular, $\ell(u_2^{-1}p_2)$ is counter-regular, and $\ell(u_1vu_2)$ is regular or counter-regular. If $\ell(u_1vu_2)$ is regular, then $\ell(p_1u_1v)$ is regular. If $\ell(u_1vu_2)$ is counter-regular, then the reduced form of $\ell(u_1vp_2)$ is counter-regular. The cycle $\langle p_2p_1u_1v \rangle$ is the contour of a




FIGURE 8. Cases 13–16, Lemma 5.10.

disc subdiagram of Δ containing the face Π_1 but not containing the face Π_2 . This contradicts the minimality of Δ .


Case 14: u_1 is a subpath of b_2^{-1} , and u_2 is a subpath of $s'_{1+}{}^{-1}$. Contradiction is obtained as in Case 13.

Case 15: u_1 is a subpath of $b_2'^{-1}$, and u_2 is a subpath of $s'_{1-}{}^{-1}$. Let p_1 and p_2 be the paths such that $u_1^{-1}p_1$ is a terminal subpath of b_2' , and p_2u_2 is an initial subpath of $y_-s'_{1-}{}^{-1}s'_{1-}{}^{-1}$. The cyclic path $p_1p_2v^{-1}u_1^{-1}$ is cyclically reduced. Suppose it is not simple. Then at least one of the following two subcases takes place:

- (a) there is a subpath of v or p_1^{-1} which represents the contour of a proper simple disc subdiagram of Δ , or
- (b) there are paths v' and p_2' such that $\langle v'p_2'^{-1} \rangle$ is the contour of a proper simple disc subdiagram of Δ , and v' and p_2' are terminal subpaths of v and p_2 , respectively.

Lemma 5.4 yields a contradiction in both subcases. Hence, $p_1p_2v^{-1}u_1^{-1}$ is a simple path. Therefore, it represents the contour of a disc subdiagram of Δ containing the faces Π_1 and Π_2 . The label of $u_1^{-1}p_1$ is counter-regular, the label of p_2u_2 is regular, and the label of $u_2^{-1}v^{-1}u_1^{-1}$ is regular or counter-regular. If $\ell(u_2^{-1}v^{-1}u_1^{-1})$ is regular, then the reduced form of $\ell(p_2v^{-1}u_1^{-1})$ is regular. If $\ell(u_2^{-1}v^{-1}u_1^{-1})$ is counter-regular, then $\ell(v^{-1}u_1^{-1}p_1)$ is counter-regular. By Lemma 5.4, $\langle p_1p_2v^{-1}u_1^{-1} \rangle = \partial\Delta$. Therefore, u_1vu_2 is a subpath of $(s'_{1-}q_1)^{-1}$. This gives a contradiction.

Case 16: u_1 is a subpath of $s'_{1+}{}^{-1}$, and u_2 is a subpath of $b_2'^{-1}$. Arguing as in Case 15, one obtains that u_1vu_2 is a subpath of $(q_1s'_{1+})^{-1}$. This gives a

FIGURE 9. The face Π in Δ and the subdiagram Δ' , Lemma 5.11.

contradiction.

A contradiction is obtained in each of the considered cases, and no other case is possible. Thus, s_2 and $(s'_{1-}q_1s'_{1+})^{-1}$ have exactly one maximal common nontrivial subpath. $\square$

Lemma 5.11. *If Π is a face of Δ , the rank of every other face of Δ is less than the rank of Π , s is the maximal selected subpath of $\partial\Pi$, and e_1 and e_n are, respectively, the initial and the terminal subpaths of s of length $|s|/n$, then all oriented edges of e_1 or all oriented edges of e_n are internal in Δ .*

Proof. Observe that proving that all oriented edges of e_1 or all oriented edges of e_n are internal in Δ is equivalent to proving that all oriented edges of e_1^{-1} or all oriented edges of e_n^{-1} are internal in the mirror copy of Δ .

Suppose Π , s , e_1 , e_n are such as in the hypotheses of the lemma. Assume that $\ell(s)$ is regular (if it is not, pass from Δ , s , e_1 , e_n to the mirror copy of Δ , s^{-1} , e_1^{-1} , e_n^{-1} , respectively).

Suppose that some oriented edge of e_1 and some oriented edge of e_n are external in Δ (see Fig. 9).

Let s' be the maximal selected external oriented arc of Δ that is a subpath of s . Let s'_- and s'_+ be such paths that $s = s'_-s's'_+$. Then s'_- is a proper initial subpath of e_1 , and s'_+ is a proper terminal subpath of e_n . Let v_1 and v_2 be the initial and the terminal subpaths of s' such that $s'_-v_1 = e_1$ and $v_2s'_+ = e_n$. Let t be the path such that $\langle st \rangle = \partial\Pi$. Let q be the path such that $\langle qs'^{-1} \rangle = \partial\Delta$. Let Δ' be the disc subdiagram of Δ obtained by removing the face Π and all edges that lie on s' together with all intermediate vertices of s' . The contour of Δ' is $\langle s'_-ts'_+q \rangle$.

Let $j = \text{rank}(\Pi)$. Then $\ell(e_1) = x_1^{m_j}$ and $\ell(e_n) = x_n^{m_j}$. The label of the first oriented edge of s' is x_1 ; the label of the last oriented edge of s' is x_n . Therefore, s' is a subpath of b_2^{-1} . Note that v_1^{-1} is an initial subpath of l_1 , v_2^{-1} is a terminal subpath of l_2 , and $v_1^{-1}qv_2^{-1} = b_1$. The reduced form of $\ell(s'_-qs'_+)$ is regular since $\ell(q)$ is regular, $\ell(s'_-)$ is a power of x_1 , and $\ell(s'_+)$ is a power of x_n . As it has been assumed, $\text{rank}(\Pi') < j$ for every face Π' of Δ' . Hence, the group word $w_j = \ell(t^{-1})$ equals a regular word (the reduced form of $\ell(s'_-qs'_+)$) modulo the relations $r = 1$, $r \in \mathcal{R}_{j-1}$. This contradicts the choice of w_i . $\square$

Lemma 5.12. *There are two distinct faces Π_1 and Π_2 , and paths s_1 , s'_1 , s'_{1-} , s'_{1+} , s_2 , s'_2 , s'_{2-} , s'_{2+} , z_1 , z_2 in Δ such that:*

- s_1 and s_2 are the maximal selected subpaths of $\partial\Pi_1$ and $\partial\Pi_2$, respectively;
- $\ell(s_1)$ is counter-regular, $\ell(s_2)$ is regular;
- $s_i = s'_{i-}s'_is'_{i+}$ for both $i = 1$ and $i = 2$;
- both s'_1 and s'_2 are maximal selected external oriented arcs of Δ ;
- $\langle z_1s'^{-1}_1z_2s'^{-1}_2 \rangle = \partial\Delta$;
- all basic letters of $\ell(s'_{2-}z_1s'_{1+})$ are in $\{x_1, \dots, x_{21}\}$, and all basic letters of $\ell(s'_{1-}z_2s'_{2+})$ are in $\{x_{n-20}, \dots, x_n\}$.

Proof. Observe that this lemma is equivalent to the analogous lemma about the mirror copy of Δ , i.e., to the statement obtained from this lemma by substituting “the mirror copy of Δ ” for “ Δ .”

Let Π_1 be a face of Δ such as in Lemma 5.9. Let s_1 be the maximal selected subpath of $\partial\Pi_1$. If $\ell(s_1)$ is regular, pass from Δ and s_1 to the mirror copy of Δ and s_1^{-1} , respectively. Hence, $\ell(s_1)$ shall be assumed to be counter-regular. Let s'_1 be the maximal selected external oriented arc of Δ that is a subpath of s_1 . Let s'_{1-} and s'_{1+} be the paths such that $s_1 = s'_{1-}s'_1s'_{1+}$. Since all of the basic letters $x_{21}, x_{22}, \dots, x_{n-20}$ occur in $\ell(s'_1)$, it follows that all basic letters of $\ell(s'_{1-})$ are in $\{x_{n-20}, \dots, x_n\}$, and all basic letters of $\ell(s'_{1+})$ are in $\{x_1, \dots, x_{21}\}$.

Let t_1 be the path such that $\langle s_1t_1 \rangle = \partial\Pi_1$. Let q_1 be the path such that $\langle s'^{-1}_1q_1 \rangle = \partial\Delta$. Let Δ' be the disc S-subdiagram of Δ obtained by removing the face Π_1 and all edges that lie on s'_1 together with all intermediate vertices of s'_1 . The contour of Δ' is $\langle s'_{1+}t_1s'_{1-}q_1 \rangle$.

It shall be shown that there is a face Π of Δ' such that Π is incident to no fewer than $(1 - 21/n)|\partial\Pi|$ selected external edges of Δ' that do not lie on t_1 . Consider 2 cases:

Case 1: there is a face Π in Δ' such that $\text{rank}(\Pi) \geq \text{rank}(\Pi_1)$. Then let Φ be the maximal simple disc S-subdiagram of Δ' that contains such a face Π . Let Σ be the sum of the degrees of all the faces of Φ . Then $\Sigma \geq |\partial\Pi| \geq |\partial\Pi_1|$. By Lemma 5.3, the number of selected external edges of Φ is at least $(1 - 2\mu)\Sigma$. Therefore, the number of selected external edges of Φ that do not lie on t_1 is at least

$$(1 - 2\mu)\Sigma - \lambda_1|\partial\Pi_1| \geq (1 - 2\mu - \lambda_1)\Sigma > \frac{n - 21}{n}\Sigma.$$

Case 2: the rank of every face of Δ' is less than the rank of Π_1 . Let e_{11} and e_{1n} be respectively the initial and terminal subpaths of s_1 of length $|s_1|/n$. Then $\ell(e_{11}) = x_n^{-m_{j_1}}$, and $\ell(e_{1n}) = x_1^{-m_{j_1}}$. Since the rank of Π_1 is greater than the rank of every other face of Δ , all oriented edges of e_{11} or all oriented edges of e_{1n} are internal in Δ by Lemma 5.11. Let e denote e_{11} in the first case or e_{1n} in the second.

If an edge lies on e and is not incident to any face of Δ' , then it also lies on t_1 . Indeed, both (mutually inverse) oriented edges corresponding to such an edge must occur in $\partial\Pi_1$; one of them is an oriented edge of e , and consequently the other cannot be an oriented edge of s_1 (two mutually inverse group letters cannot both occur in $\ell(s_1)$) and has to be an oriented edge of t_1 . Since

$$|t_1| = |w_{j_1}| < m_{j_1} = |e|$$

(see (5.5)), the ratio of the number of edges that lie on t_1 and are incident with faces of Δ' to the number of edges that lie on e but are also incident with faces of Δ' is not greater than

$$\frac{|t_1|}{|e|} = \frac{|w_{j_1}|}{m_{j_1}} \leq \frac{n\lambda_1}{1-\lambda_1}$$

(see (5.4)). Let Φ be a maximal simple disc S-subdiagram of Δ' such that the ratio of the number of its (external) edges that lie on t_1 to the number of its (external) edges that lie on e is not greater than

$$\frac{n\lambda_1}{1-\lambda_1}.$$

Let Σ be the sum of the degrees of all the faces of Φ . Since all edges that lie on e are labelled with a same basic letter, the number of edges of Φ that lie on e is less than

$$\left(\lambda_1 + \frac{1-\lambda_1}{n}\right)\Sigma$$

(because the number of edges with a same label incident to a given face Π of Δ is less than $(\lambda_1 + (1-\lambda_1)/n)|\partial\Pi|$). By Lemma 5.3, the number of selected external edges of Φ is at least $(1-2\mu)\Sigma$. Therefore, the number of selected external edges of Φ that do not lie on t_1 is greater than

$$\left(1-2\mu - \frac{n\lambda_1}{1-\lambda_1}\left(\lambda_1 + \frac{1-\lambda_1}{n}\right)\right)\Sigma = \left(1-2\mu - \lambda_1 - \frac{n\lambda_1^2}{1-\lambda_1}\right)\Sigma > \frac{n-21}{n}\Sigma$$

(see (5.2)).

In both cases, there is a face Π in Φ such that Π is incident to at least $(1-21/n)|\partial\Pi|$ selected external edges of Φ (of Δ') that do not lie on t_1 . Let Π_2 be such a face. Let s_2 be the maximal selected subpath of $\partial\Pi_2$. Then the paths s_2 and $(s'_{1-}q_1s'_{1+})^{-1}$ have at least $(1-21/n)|\partial\Pi_2|$ common oriented edges. Let $\tilde{s}_2$ be the maximal common nontrivial subpath of s_2 and $(s'_{1-}q_1s'_{1+})^{-1}$ (see Lemma 5.10). Note that $\tilde{s}_2$ is a selected external oriented arc of Δ' . Every selected external edge of Δ' incident to Π_2 lies either on t_1 or on $\tilde{s}_2$. Therefore,

$$|\tilde{s}_2| \geq \frac{n-21}{n}|\partial\Pi_2| > \frac{n-21}{n}|s_2|.$$

Therefore, $\ell(\tilde{s}_2)$ has at least $n-20$ distinct basic letters, which implies that each of the basic letters $x_{21}, \dots, x_{n-20}$ occurs in it. The basic letters $x_{22}, \dots, x_{n-21}$ do not occur on s'_{1-} nor on s'_{1+} , but they occur in $\ell(\tilde{s}_2)$. Hence, the paths $\tilde{s}_2$ and q_1^{-1} have common oriented edges. Therefore, Π_2 is incident to some selected external edges of Δ .

Let s'_2 be the maximal selected external oriented arc of Δ that is a subpath of s_2 . Then s'_2 is the maximal common subpath of $\tilde{s}_2$ and q_1^{-1} . All of the basic letters $x_{22}, \dots, x_{n-21}$ occur in $\ell(s'_2)$ (since they occur in $\ell(\tilde{s}_2)$ but neither in $\ell(s'_{1-})$ nor in $\ell(s'_{1+})$).

Since $\ell(s_1)$ is counter-regular, s'_1 is a subpath of b_1^{-1} . Therefore, s'_2 is a subpath of b_2^{-1} , and $\ell(s_2)$ is regular.

Let s'_{2-} and s'_{2+} be the paths such that $s_2 = s'_{2-}s'_2s'_{2+}$. Let z_1 and z_2 be the paths such that $\langle z_1s'^{-1}_1z_2s'^{-1}_2 \rangle = \partial\Delta$. It shall be proved that all basic

letters of $\ell(s'_{2-})$ and $\ell(z_1)$ are among $x_1, \dots, x_{21}$, and all basic letters of $\ell(s'_{2+})$ and $\ell(z_2)$ are among $x_{n-20}, \dots, x_n$.

First, consider s'_{2-} and z_1 .

Case 1: $\tilde{s}_2$ has no common oriented edges with s'^{-1}_{1+} . Then s'_2 is an initial subpath of $\tilde{s}_2$. Therefore, the basic letter x_{21} , along with $x_{22}, \dots, x_{n-21}$, occurs in $\ell(s'_2)$. Hence, all basic letters of $\ell(s'_{2-})$ are in $\{x_1, \dots, x_{21}\}$. The label of every edge that lies on both z_1 and b_1 is in $\{x_1, \dots, x_{21}\}$ since the basic letter x_{21} occurs in $\ell(s'_1)$, and the only maximal common subpath of $z_1 s'^{-1}_1$ and b_1 is an initial subpath of b_1 . The label of every edge that lies on both z_1 and b_2 is in $\{x_1, \dots, x_{21}\}$ since the basic letter x_{21} occurs in $\ell(s'_2)$, and the only maximal common subpath of $s'^{-1}_2 z_1$ and b_2 is a terminal subpath of b_2 .

Case 2: $\tilde{s}_2$ has at least one oriented edge in common with s'^{-1}_{1+} . Then some nontrivial terminal subpath of s'^{-1}_{1+} is an initial subpath of $\tilde{s}_2$, and z_1 is trivial. The terminal subpath of s'^{-1}_{1+} that is an initial subpath of $\tilde{s}_2$ is also a terminal subpath of s'_{2-} . Since s'^{-1}_{1+} and s'_{2-} have a common nontrivial terminal subpath, the sets of basic letters of their labels coincide. Therefore, all basic letters of $\ell(s'_{2-})$ are in $\{x_1, \dots, x_{21}\}$. The label of z_1 is empty.

Second, consider s'_{2+} and z_2 in the same manner.

Case 1: $\tilde{s}_2$ has no common oriented edges with s'^{-1}_{1-} . Then s'_2 is a terminal subpath of $\tilde{s}_2$. Therefore, the basic letter x_{n-20} , along with $x_{22}, \dots, x_{n-21}$, occurs in $\ell(s'_2)$. Hence, all basic letters of $\ell(s'_{2+})$ are in $\{x_{n-20}, \dots, x_n\}$. The label of every edge that lies on both z_2 and b_1 is in $\{x_{n-20}, \dots, x_n\}$ since the basic letter x_{n-20} occurs in $\ell(s'_1)$, and the only maximal common subpath of $s'^{-1}_1 z_2$ and b_1 is a terminal subpath of b_1 . The label of every edge that lies on both z_2 and b_2 is in $\{x_{n-20}, \dots, x_n\}$ since the basic letter x_{n-20} occurs in $\ell(s'_2)$, and the only maximal common subpath of $z_2 s'^{-1}_2$ and b_2 is an initial subpath of b_2 .

Case 2: $\tilde{s}_2$ has at least one oriented edge in common with s'^{-1}_{1-} . Then some nontrivial initial subpath of s'^{-1}_{1-} is a terminal subpath of $\tilde{s}_2$, and z_2 is trivial. The initial subpath of s'^{-1}_{1-} that is a terminal subpath of $\tilde{s}_2$ is also an initial subpath of s'_{2+} . Since s'^{-1}_{1-} and s'_{2+} have a common nontrivial initial subpath, the sets of basic letters of their labels coincide. Therefore, all basic letters of $\ell(s'_{2+})$ are in $\{x_{n-20}, \dots, x_n\}$. The label of z_2 is empty.

Clearly, the faces Π_1 , Π_2 and the paths s_1 , s'_1 , s'_{1-} , s'_{1+} , s_2 , s'_2 , s'_{2-} , s'_{2+} , z_1 , z_2 are desired ones. $\square$

Lemma 5.13. *The diagram Δ has more than 2 faces.*

Proof. Note that Δ obviously has the same number of faces as its mirror copy.

Suppose the statement is not true, i.e., Δ has no more than 2 faces. (Then, by Lemma 5.5, it has exactly 2 faces.)

Let Π_1 and Π_2 be such faces of Δ as in Lemma 5.12. Then the diagram Δ consists of the faces Π_1 and Π_2 attached to each other along a common arc. If $\text{rank}(\Pi_2) > \text{rank}(\Pi_1)$, pass to the mirror copy of Δ and interchange the roles of Π_1 and Π_2 . Hence, it shall be assumed that $\text{rank}(\Pi_1) \geq \text{rank}(\Pi_2)$.

Let $s_1, s'_1, s'_{1-}, s'_{1+}, s_2, s'_2, s'_{2-}, s'_{2+}$ be the subpaths of $\partial\Pi_1$ and $\partial\Pi_2$ such as in Lemma 5.12. Then all basic letters of the labels of $s'_{1-}, s'_{1+}, s'_{2-}, s'_{2+}$ are in $\{x_1, \dots, x_{21}\} \cup \{x_{n-20}, \dots, x_n\}$, and s'_i is a subpath of b_i^{-1} for $i = 1$ and $i = 2$.

There are exactly two (mutually inverse) maximal internal oriented arcs in Δ . Denote the one that is a subpath of $\partial\Pi_1$ by u .

Let t_1 and t_2 be the path such that $\langle s_1 t_1 \rangle = \partial\Pi_1$ and $\langle s_2 t_2 \rangle = \partial\Pi_2$. Each of the faces Π_1 and Π_2 is incident to exactly one maximal selected external arc of Δ . Therefore, each of the paths t_1 and t_2 has a common vertex with u .

Suppose t_1 and t_2 do not have common vertices. Then either some (proper) nontrivial initial subpath of s_1 is inverse to a (proper) initial subpath of s_2 , or some (proper) nontrivial terminal subpath of s_1 is inverse to a (proper) terminal subpath of s_2 . Both cases are impossible since no proper nontrivial prefix of $\ell(s_2)$ can be a suffix of $\ell(s_1^{-1})$, and no proper nontrivial suffix of $\ell(s_2)$ can be a prefix of $\ell(s_1^{-1})$. (A contradiction may also be obtained by applying Lemma 5.10.) Hence, the paths t_1 and t_2 have at least one common vertex.

Let $j_1 = \text{rank}(\Pi_1)$ and $j_2 = \text{rank}(\Pi_2)$. Let e_{i1} and e_{in} be respectively the initial and the terminal subpaths of s_i of length $m_{j_i} = |s_i|/n$ for $i = 1$ and $i = 2$. Then $\ell(e_{11}) = x_n^{-m_{j_1}}$, $\ell(e_{1n}) = x_1^{-m_{j_1}}$, $\ell(e_{21}) = x_1^{m_{j_2}}$, and $\ell(e_{2n}) = x_n^{m_{j_2}}$. At most one of the two group letters $x_1^{\pm 1}$ and at most one of $x_n^{\pm 1}$ can occur as labels of oriented edges of $\partial\Delta$. Therefore, at least one of the paths e_{1n} and e_{21} and at least one of the paths e_{11} and e_{2n} do not have external oriented edges in Δ .

Suppose $\text{rank}(\Pi_1) = \text{rank}(\Pi_2)$. Then

$$|e_{11}| = |e_{1n}| = |e_{21}| = |e_{2n}| = m_{j_1} \geq |w_{j_1}| = |t_1| = |t_2|.$$

Recall that w_{j_1} does not start with $x_1^{\pm 1}$, does not end with $x_n^{\pm 1}$, and is not a letter power. Therefore, every subpath of either $\partial\Pi_1$ or $\partial\Pi_2$ labelled with a letter power has length of at most m_{j_1} . Moreover, for every i , there exist exactly one subpath of $\partial\Pi_1$ and exactly one subpath of $\partial\Pi_2$ of length m_{j_1} labelled with powers of x_i (i.e., with $x_i^{\pm m_{j_1}}$). If e_{1n} has no external oriented edges in Δ , then it is a subpath of u . If e_{21} has no external oriented edges in Δ , then it is a subpath of u^{-1} . In either case u has a subpath labelled with $x_1^{-m_{j_1}}$, and the pair $\{\Pi_1, \Pi_2\}$ is immediately cancellable. This contradicts Lemma 5.1. Hence, $\text{rank}(\Pi_1) > \text{rank}(\Pi_2)$.

Let Δ' be the disc S-subdiagram of Δ obtained by removing the face Π_1 and all edges and intermediate vertices of s'_1 . The only face of Δ' is Π_2 .

By Lemma 5.11, all oriented edges of e_{11} or all oriented edges of e_{1n} are internal in Δ . Consider the following 5 cases (see Fig. 10):

(Notice that Case 3 is “symmetric” with Case 2, and Case 5—with Case 4.)

Case 1: each of the paths e_{11}, e_{1n}, e_{21} , and e_{2n} has an internal oriented edge. Then the paths s'_{1-} and s'_{2+} have a common nontrivial terminal subpath, the paths s'_{1+} and s'_{2-} have a common nontrivial initial subpath, and $\partial\Delta = \langle s'^{-1}_{1-} s'^{-1}_{2-} \rangle$. Let v_1 be a common nontrivial terminal subpath of s'_{1-} and s'^{-1}_{2-} , and let v_2 be a common nontrivial initial subpath of s'_{1+} and s'^{-1}_{2-} . The contour of Δ' is $\langle s'_{1+} t_1 s'_{1-} s'^{-1}_{2-} \rangle$. The label of $s'_{1-} s'^{-1}_{2-} s'_{1+}$ is





FIGURE 10. Cases 1–5, Lemma 5.13.

counter-regular because the labels of s'_{1-} , $v_1 s_2'^{-1} v_2$, and s'_{1+} are such. Hence, the group word $w_{j_1} = \ell(t_1)$ equals the regular word $\ell(s'_{1-} s_2'^{-1} s'_{1+})^{-1}$ modulo the relation $r_{j_2} = 1$. This contradicts the choice of w_{j_1} .

Case 2: all oriented edges of e_{1n} are external (consequently, all oriented edges of e_{11} and e_{21} are internal), and some oriented edge of e_{2n} is internal. Then the paths s'_{1-} and $s_2'^{-1}$ have a common nontrivial terminal subpath. Let v_1 be a common nontrivial terminal subpath of s'_{1-} and $s_2'^{-1}$. Let z_1 be the path such that $\langle z_1 s_1'^{-1} s_2'^{-1} \rangle = \partial\Delta$. Then z_1^{-1} is an initial subpath of t_1 . Since $\ell(e_{1n})$ is a power of x_1 , the path e_{1n}^{-1} is a (terminal) subpath of b_2 . Therefore, $s_2'^{-1} z_1 e_{1n}^{-1}$ is a subpath of b_2 . The contour of Δ' is $\langle z_1 t_1 s'_{1-} s_2'^{-1} \rangle$. The label of $s'_{1-} s_2'^{-1} z_1$ is counter-regular because the labels of s'_{1-} , $v_1 s_2'^{-1}$, and $s_2'^{-1} z_1$ are such. Hence, the group word $w_{j_1} = \ell(t_1)$ equals the regular word $\ell(s'_{1-} s_2'^{-1} z_1)^{-1}$ modulo the relation $r_{j_2} = 1$. This contradicts the choice of w_{j_1} .

Case 3: all oriented edges of e_{11} are external (consequently, all oriented

FIGURE 11. The subdiagram Δ' , Lemma 5.14.

edges of e_{1n} and e_{2n} are internal), and some oriented edge of e_{21} is internal. Then the paths s'_{1+} and s'^{-1}_{2-} have a common nontrivial initial subpath. Let v_2 be a common nontrivial initial subpath of s'_{1+} and s'^{-1}_{2-} . Let z_2 be the path such that $\langle s'^{-1}_1 z_2 s'^{-1}_2 \rangle = \partial\Delta$. Then z_2^{-1} is a terminal subpath of t_1 . Since $\ell(e_{11})$ is a power of x_n , the path e_{11}^{-1} is an (initial) subpath of b_2 . Therefore, $e_{11}^{-1} z_2 s'^{-1}_2$ is a subpath of b_2 . The contour of Δ' is $\langle s'_{1+} t_1 z_2 s'^{-1}_2 \rangle$. The label of $z_2 s'^{-1}_2 s'_{1+}$ is counter-regular because the labels of $z_2 s'^{-1}_2$, $s'^{-1}_2 v_2$, and s'_{1+} are such. Hence, the group word $w_{j_1} = \ell(t_1)$ equals the regular word $\ell(z_2 s'^{-1}_2 s'_{1+})^{-1}$ modulo the relation $r_{j_2} = 1$. This contradicts the choice of w_{j_1} .

Case 4: all oriented edges of e_{21} are external. Consequently, all oriented edges of e_{1n} are internal. Then e_{1n} is a terminal subpath of s'_{1+} . The path s'^{-1}_{1+} is a subpath of t_2 (because t_2 has common vertices with both e_{21} and t_1). Therefore, $|t_2| \geq |s'_{1+}| \geq |e_{1n}| = m_{j_1} > |w_{j_2}| = |t_2|$ (see (5.5)). This gives a contradiction.

Case 5: all oriented edges of e_{2n} are external. Consequently, all oriented edges of e_{11} are internal. Then e_{11} is an initial subpath of s'_{1-} . The path s'^{-1}_{1-} is a subpath of t_2 (because t_2 has common vertices with both e_{2n} and t_1). Therefore, $|t_2| \geq |s'_{1-}| \geq |e_{11}| = m_{j_1} > |w_{j_2}| = |t_2|$ (see (5.5)). This gives a contradiction.

No other case is possible. Thus, Δ has more than 2 faces. $\square$

Lemma 5.14. *Let Π_1 and Π_2 be distinct faces of Δ such as in Lemma 5.12. Then the ranks of Π_1 and Π_2 are distinct.*

Proof. Let Π_1 , Π_2 , s_1 , s'_1 , s'_{1-} , s'_{1+} , s_2 , s'_2 , s'_{2-} , s'_{2+} , z_1 , z_2 be as in Lemma 5.12. Let $c_1 = s'_{2-} z_1 s'_{1+}$ and $c_2 = s'_{1-} z_2 s'_{2+}$. Let t_1 and t_2 be the paths such that $\langle s_1 t_1 \rangle = \partial\Pi_1$ and $\langle s_2 t_2 \rangle = \partial\Pi_2$. Let Δ' be the disc S-subdiagram of Δ obtained by removing the faces Π_1 and Π_2 and all edges and intermediate vertices of the paths s'_1 and s'_2 (see Fig. 11).

The contour of Δ' is $\langle c_1 t_1 c_2 t_2 \rangle$. By Lemma 5.13, the diagram Δ' is non-degenerate. Therefore, by Lemma 5.4, the label of $c_1 t_1 c_2 t_2$ is not freely trivial.

Suppose $\text{rank}(\Pi_1) = \text{rank}(\Pi_2)$. Let $j = \text{rank}(\Pi_1) = \text{rank}(\Pi_2)$. Recall that $\ell(t_1) = w_j$, $\ell(t_2) = w_j^{-1}$, all basic letters of $\ell(c_1)$ are in $\{x_1, \dots, x_{21}\}$,

and all basic letters of $\ell(c_2)$ are in $\{x_{n-20}, \dots, x_n\}$. Let $h_1 = \ell(c_1^{-1})$ and $h_2 = \ell(c_2)$. Then $[h_1] = [w_j][h_2][w_j]^{-1}$. Let F be the subgroup of G generated by $\{a_1, \dots, a_{21}\} \cup \{a_{n-20}, \dots, a_n\}$. By Property 1, the group F is free of rank 42 (since $42 \leq n - 21$), and elements of F are conjugate in G if and only if they are conjugate in F . The elements $[h_1]$ and $[h_2]$ are in F . Moreover, $[h_1]$ belongs to the subgroup generated by $\{a_1, \dots, a_{21}\}$, and $[h_2]$ belongs to the subgroup generated by $\{a_{n-20}, \dots, a_n\}$. Since $[h_1]$ and $[h_2]$ are conjugate in G , they are conjugate in F and therefore $[h_1] = [h_2] = 1$. Hence, $\ell(c_1 t_1 c_2 t_2)$ is freely trivial. This gives a contradiction. $\square$

Lemma 5.15. *Let Π_1 and Π_2 be distinct faces of Δ , both incident to selected external edges of Δ . Suppose the rank of Π_1 is greater than the rank of every other face of Δ . Let s_1 and s_2 be the maximal selected subpaths of $\partial\Pi_1$ and $\partial\Pi_2$, respectively. Suppose the label of one of the paths s_1 or s_2 is regular, and the label of the other is counter-regular. Let s'_1 and s'_2 be the maximal selected external oriented arcs of Δ that are subpaths of s_1 and s_2 , respectively. Suppose there are at least 2 distinct basic letters in the label of each of the paths s'_1 and s'_2 . Let e_{11} and e_{1n} be respectively the initial and the terminal subpaths of s_1 of length $|s_1|/n$. Then at least one of the paths e_{11} or e_{1n} has the property that every edge that lies on it is internal in Δ and does not lie on s_2 .*

Proof. Observe that it suffices to prove that at least one of the paths e_{1n}^{-1} or e_{11}^{-1} has the property that every edge that lies on it is internal in the mirror copy of Δ and does not lie on s_2^{-1} .

Assume that $\ell(s_1)$ is counter-regular (if it is not, pass from Δ , b_1 , b_2 , l_1 , l_2 , s_1 , s_2 , s'_1 , s'_2 , e_{11} , e_{1n} to the mirror copy of Δ , b_2^{-1} , b_1^{-1} , l_1^{-1} , l_2^{-1} , s_1^{-1} , s_2^{-1} , $s_1'^{-1}$, $s_2'^{-1}$, e_{11}^{-1} , e_{1n}^{-1} , respectively). Then s'_1 is a subpath of b_1^{-1} , and s'_2 is a subpath of b_2^{-1} .

Let s'_{1-} and s'_{1+} be the paths such that $s_1 = s'_{1-} s'_1 s'_{1+}$. Let t_1 and t_2 be the paths such that $\langle s_1 t_1 \rangle = \partial\Pi_1$ and $\langle s_2 t_2 \rangle = \partial\Pi_2$. Let q_1 be the path such that $\langle s_1'^{-1} q_1 \rangle = \partial\Delta$. Let Δ' be the disc subdiagram of Δ obtained by removing the face Π_1 and all edges that lie on s'_1 together with all intermediate vertices of s'_1 . The contour of Δ' is $\langle s'_{1+} t_1 s'_{1-} q_1 \rangle$. Let e_{21} and e_{2n} be respectively the initial and terminal subpaths of s_2 of length $|s_2|/n$.

Denote the ranks of Π_1 and Π_2 by j_1 and j_2 , respectively. Then

$$\begin{aligned} \ell(s_1) &= x_n^{-m_{j_1}} x_{n-1}^{-m_{j_1}} \dots x_1^{-m_{j_1}}, & \ell(t_1) &= w_{j_1}, \\ \ell(s_2) &= x_1^{m_{j_2}} x_2^{m_{j_2}} \dots x_n^{m_{j_2}}, & \ell(t_2) &= w_{j_2}^{-1}, \\ \ell(e_{11}) &= x_n^{-m_{j_1}}, & \ell(e_{1n}) &= x_1^{-m_{j_1}}, & \ell(e_{21}) &= x_1^{m_{j_2}}, & \ell(e_{2n}) &= x_n^{m_{j_2}}. \end{aligned}$$

At most one of the two group letters $x_1^{\pm 1}$ and at most one of $x_n^{\pm 1}$ can occur as labels of oriented edges of $\partial\Delta$ (the labels of the representatives of $\partial\Delta$ are cyclically reduced). Therefore, at least one of the paths e_{1n} or e_{21} and at least one of the paths e_{11} or e_{2n} do not have external oriented edges in Δ . By Lemma 5.11, all oriented edges of e_{11} or all oriented edges of e_{1n} are internal in Δ .

Suppose the conclusion of the lemma does not hold. Then some oriented edge of e_{11} is either external in Δ or inverse to an oriented edge of e_{2n} , and




FIGURE 12. Cases 1–3, Lemma 5.15.

some oriented edge of e_{1n} is either external in Δ or inverse to an oriented edge of e_{21} . Therefore, as follows from Lemma 5.11, at least one of the following 3 cases takes place:

- some oriented edge of e_{11} is inverse to an oriented edge of e_{2n} , and some oriented edge of e_{1n} is inverse to an oriented edge of e_{21} , or
- some oriented edge of e_{11} is external in Δ , and some oriented edge of e_{1n} is inverse to an oriented edge of e_{21} , or
- some oriented edge of e_{11} is inverse to an oriented edge of e_{2n} , and some oriented edge of e_{1n} is external in Δ .

In every case the path s_2 has a common oriented edge with s'_{1-} or s'_{1+} . Let $\tilde{s}_2$ be the maximal common nontrivial subpath of s_2 and $(s'_{1-}q_1s'_{1+})^{-1}$ (see Lemma 5.10). Now, consider each of the 3 cases separately (see Fig. 12).

(Notice that Case 3 is “symmetric” with Case 2.)

Case 1: some oriented edge of e_{11} is inverse to an oriented edge of e_{2n} , and some oriented edge of e_{1n} is inverse to an oriented edge of e_{21} . Then some nontrivial initial subpath of $\tilde{s}_2^{-1}$ is a terminal subpath of s'_{1-} , and some nontrivial terminal subpath of $\tilde{s}_2^{-1}$ is an initial subpath of s'_{1+} (because $\tilde{s}_2^{-1}$ is a subpath of $s'_{1-}q_1s'_{1+}$). Let v_1 and v_2 be the paths such that $v_1q_1v_2 = \tilde{s}_2^{-1}$. Then v_1 is a nontrivial terminal subpath of s'_{1-} , and v_2 is a nontrivial initial subpath of s'_{1+} . The label of $s'_{1-}q_1s'_{1+}$ is counter-regular since the labels of s'_{1-} , $v_1q_1v_2$, and s'_{1+} are such. As it is assumed, $\text{rank}(\Pi) < j_1$ for every face Π of Δ' . Recall that $\langle s'_{1+}t_1s'_{1-}q_1 \rangle = \partial\Delta'$. Hence, the group word $w_{j_1} = \ell(t_1)$ equals the regular word $\ell(s'_{1-}q_1s'_{1+})^{-1}$ modulo the relations $r = 1$, $r \in \mathcal{R}_{j_1-1}$. This contradicts the choice of w_{j_1} .

Case 2: some oriented edge of e_{11} is external in Δ , and some oriented edge of e_{1n} is inverse to an oriented edge of e_{21} . Then s'_{1-} is a proper initial subpath of e_{11} . Let v_1 be the initial subpath of s'_1 such that $s'_{1-}v_1 = e_{11}$. Then v_1^{-1} is a subpath of l_2 since $\ell(v_1^{-1})$ is a nontrivial power of x_n . Some nontrivial terminal subpath of $\tilde{s}_2^{-1}$ is an initial subpath of s'_{1+} . Therefore, s'^{-1}_2 is a terminal subpath of q_1 , and $v_1^{-1}q_1$ is a subpath of b_2 (because s'^{-1}_2 is a common subpath of q_1 and b_2). Let v_2 be the path such that $s'^{-1}_2v_2$ is a terminal subpath of $\tilde{s}_2^{-1}$. Then v_2 is a nontrivial initial subpath of s'_{1+} . The label of $q_1s'_{1+}$ is counter-regular since the labels of q_1 , $s'^{-1}_2v_2$, and s'_{1+} are such. The reduced form of $\ell(s'_{1-}q_1s'_{1+})$ is counter-regular since $\ell(s'_{1-})$ is a power of x_n , and $\ell(q_1s'_{1+})$ is counter-regular. As it is assumed, $\text{rank}(\Pi) < j_1$ for every face Π of Δ' . Hence, the group word $w_{j_1} = \ell(t_1)$ equals a regular word (the reduced form of $\ell(s'_{1-}q_1s'_{1+})^{-1}$) modulo the relations $r = 1$, $r \in \mathcal{R}_{j_1-1}$. This contradicts the choice of w_{j_1} .

Case 3: some oriented edge of e_{11} is inverse to an oriented edge of e_{2n} , and some oriented edge of e_{1n} is external in Δ . Contradiction is obtained as in Case 2.

No other case is possible. Thus, at least one of the paths e_{11} or e_{1n} has no common oriented edges with s_2^{-1} and no external oriented edges. $\square$

Lemma 5.16. *Let $\Pi_1, \Pi_2, s_1, s'_1, s'_{1-}, s'_{1+}, s_2, s'_2, s'_{2-}, s'_{2+}, z_1, z_2$ be such as in Lemma 5.12. Let $c_1 = s'_{2-}z_1s'_{1+}$ and $c_2 = s'_{1-}z_2s'_{2+}$. Then Δ has a simple disc S-subdiagram Φ such that Π_1 and Π_2 are not in Φ , and the total number of selected external edges of Φ that lie on c_1 or c_2 is greater than*

$$\frac{n-21}{n} \sum_{\Pi \in \Phi(2)} |\partial\Pi|.$$

Proof. Observe that to prove this lemma, it suffices to prove the analogous lemma for the mirror copy of Δ .

By Lemma 5.14, $\text{rank}(\Pi_1) \neq \text{rank}(\Pi_2)$. If $\text{rank}(\Pi_1) < \text{rank}(\Pi_2)$, pass from $\Delta, \Pi_1, \Pi_2, s_1, s'_1, s'_{1-}, s'_{1+}, s_2, s'_2, s'_{2-}, s'_{2+}, z_1, z_2, c_1, c_2$ to the mirror copy of $\Delta, \Pi_2, \Pi_1, s_2^{-1}, s'_{2+}{}^{-1}, s'^{-1}_2, s'_{2-}{}^{-1}, s_1^{-1}, s'_{1+}{}^{-1}, s'^{-1}_1, s'_{1-}{}^{-1}, z_1^{-1}, z_2^{-1}, c_1^{-1}, c_2^{-1}$, respectively. Hence, assume that $j_1 = \text{rank}(\Pi_1) > \text{rank}(\Pi_2) = j_2$.

Let t_1 and t_2 be the paths such that $\langle s_1t_1 \rangle = \partial\Pi_1$ and $\langle s_2t_2 \rangle = \partial\Pi_2$. Let Δ' be the disc S-subdiagram of Δ obtained by removing the faces Π_1 and Π_2 and all edges that lie on s'_1 or s'_2 together with all intermediate vertices of s'_1 and s'_2 . The contour of Δ' is $\langle c_1t_1c_2t_2 \rangle$. By Lemma 5.13, the diagram Δ' is nondegenerate.

Consider 2 cases:

Case 1: there is a face Π in Δ' such that $\text{rank}(\Pi) \geq \text{rank}(\Pi_1)$. Then let Φ be the maximal simple disc S-subdiagram of Δ' that contains such a face Π . Let Σ be the sum of the degrees of all the faces of Φ . Then $\Sigma \geq |\partial\Pi| \geq |\partial\Pi_1| > |\partial\Pi_2|$. By Lemma 5.3, the number of selected external edges of Φ is at least $(1-2\mu)\Sigma$. Therefore, the number of selected external edges of Φ that do not lie on t_1 nor on t_2 is at least

$$(1-2\mu)\Sigma - \lambda_1|\partial\Pi_1| - \lambda_1|\partial\Pi_2| > (1-2\mu-2\lambda_1)\Sigma > \frac{n-21}{n}\Sigma.$$

Case 2: the rank of every face of Δ' is less than the rank of Π_1 . Let e_{11} and e_{1n} be respectively the initial and the terminal subpaths of s_1 of length $m_{j_1} = |s_1|/n$. By Lemma 5.15, at least one of the paths e_{11} or e_{1n} has the property that every edge that lies on it either is incident to a face of Δ' or lies on t_1 or t_2 . Let e be one of the paths e_{11} or e_{1n} with this property.

Since

$$|t_1| + |t_2| = |w_{j_1}| + |w_{j_2}| < \frac{1}{2}m_{j_1} + \frac{1}{2}m_{j_1} = m_{j_1} = |e|$$

(see (5.5)), the ratio of the number of edges that lie on t_1 or t_2 and are incident with faces of Δ' (consequently, do not lie on e) to the number of edges that lie on e and are incident with faces of Δ' (equivalently, lie neither on t_1 nor on t_2) is at most

$$\frac{|t_1| + |t_2|}{|e|} = \frac{|w_{j_1}| + |w_{j_2}|}{m_{j_1}} \leq \frac{2n\lambda_1}{1 - \lambda_1}$$

(see (5.4)). Let Φ be a maximal simple disc S-subdiagram of Δ' such that the ratio of the number of its (external) edges that lie on t_1 or t_2 to the number of its (external) edges that lie on e is at most

$$\frac{2n\lambda_1}{1 - \lambda_1}.$$

Let Σ be the sum of the degrees of all the faces of Φ . Since all the edges that lie on e are labelled with a same basic letter, the number of edges of Φ that lie on e is less than

$$\left(\lambda_1 + \frac{1 - \lambda_1}{n}\right)\Sigma.$$

By Lemma 5.3, the number of selected external edges of Φ is at least $(1 - 2\mu)\Sigma$. Therefore, the number of selected external edges of Φ that neither lie on t_1 nor on t_2 is greater than

$$\left(1 - 2\mu - \frac{2n\lambda_1}{1 - \lambda_1} \left(\lambda_1 + \frac{1 - \lambda_1}{n}\right)\right)\Sigma = \left(1 - 2\mu - 2\lambda_1 - \frac{2n\lambda_1^2}{1 - \lambda_1}\right)\Sigma \geq \frac{n - 21}{n}\Sigma$$

(see (5.2)).

Every external edge of Φ that neither lies on t_1 nor on t_2 lies on c_1 or c_2 . Thus, the desired inequality holds in the both cases. $\square$

Proof of Property 2. The existence is obvious from the choice of relators.

Suppose that presentation of some element of G in the form $a_1^{k_1} \dots a_n^{k_n}$ is not unique. Then there exist two distinct regular group words over $\mathfrak{A}$ whose values in G coincide. Consider such a pair of group words with the minimal sum of their lengths. Consider a minimal deduction diagram for the equality of these group words over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$. More precisely, let Δ be a special disc S-diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ whose contour has a representative of the form $p_1 p_2$ such that $\ell(p_1)$ and $\ell(p_2^{-1})$ are distinct regular group words, and $\ell(p_1 p_2)$ is cyclically reduced; moreover, let Δ be such an S-diagram with the minimal possible number of faces.

Choose faces Π_1 and Π_2 and paths $s_1, s'_1, s'_{1-}, s'_{1+}, s_2, s'_2, s'_{2-}, s'_{2+}, z_1, z_2$ as in Lemma 5.12. Denote the paths $s'_{2-} z_1 s'_{1+}$ and $s'_{1-} z_2 s'_{2+}$ by c_1 and c_2 , respectively (as in the hypotheses of Lemma 5.16). Apply Lemma 5.16:

let Φ be a simple disc S-subdiagram of Δ such that the number of selected external edges of Φ that lie on c_1 or c_2 is greater than

$$\frac{n-21}{n} \sum_{\Pi \in \Phi(2)} |\partial \Pi|.$$

By the choice of Π_1 and Π_2 , all basic letters of $\ell(c_1)$ are in $\{x_1, \dots, x_{21}\}$, and all basic letters of $\ell(c_2)$ are in $\{x_{n-20}, \dots, x_n\}$. Therefore, by Proposition 5.2, the number of selected external edges of Φ that lie on c_1 or c_2 is less than

$$\frac{42}{n} \sum_{\Pi \in \Phi(2)} |\partial \Pi|.$$

Since $n-21 \geq 42$, one has a contradiction. The uniqueness is proved. $\square$

Remark 5.1. The “symmetry” mentioned in the proofs of Lemmas 5.8, 5.10, 5.13, and 5.15 is the symmetry between the S-diagram Δ and the S-diagram obtained from the mirror copy of Δ by re-labelling according to the following rule: if the label of an oriented edge e is x_i^σ , then re-label e with $x_{n+1-i}^{-\sigma}$ (this gives a diagram not over the same presentation).

The proof of the next property uses a homological argument.

Property 3. *The group G is torsion-free.*

Proof. The group presentation $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ is strongly aspherical by Corollary 5.1.2 of Proposition 5.1; therefore, it is aspherical in the sense of [Ol’91]. Since no element of $\mathcal{R}$ represents a proper power in the free group on $\mathfrak{A}$, the relation module M of $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ is a free G -module by Corollary 32.1 in [Ol’91]. Therefore, there exists a finite-length free resolution of $\mathbb{Z}$ over $\mathbb{Z}G$:

$$(5.6) \quad 0 \rightarrow M \rightarrow \bigoplus_{x \in \mathfrak{A}} \mathbb{Z}G \rightarrow \mathbb{Z}G \rightarrow \mathbb{Z} \rightarrow 0,$$

where $\mathbb{Z}G$ and $\bigoplus_{x \in \mathfrak{A}} \mathbb{Z}G$ are identified with the G -modules of, respectively, 0- and 1-dimensional cellular chains of the Cayley complex of $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ (see [Bro94]).

Now suppose that G has torsion. Let H be a non-trivial finite cyclic subgroup of G . Every free G -module may be naturally regarded as a free H -module. Hence, the resolution (5.6) may be viewed as a free resolution of $\mathbb{Z}$ over $\mathbb{Z}H$. This contradicts the fact that all odd-dimensional homology groups of H are nontrivial. $\square$

For more results on group presentations with various forms of asphericity, and torsion in such groups, see [CCH81], [Hue79], [Hue80], and Theorems 32.1, 32.2 in [Ol’91].

Properties 1, 2, and 3 of G demonstrate that the answer to Bludov’s question is negative.

Remark 5.2. If n was chosen to be less than 63 but greater than 26 ($26 < n < 63$), then Properties 1 and 3 would still hold, but the proof of Property 2 would not work.

For every $i \in \mathbb{N} \cup \{0\}$, let G_i be the group defined by the presentation $\langle \mathfrak{A} \parallel \mathcal{R}_i \rangle$ (see Section 2). Also for every $i \in \mathbb{N} \cup \{0\}$, let ϕ_i be the natural epimorphism $G_i \rightarrow G_{i+1}$.

Property 4. *All groups $G_0, G_1, G_2, \dots$ are hyperbolic, and G is isomorphic to the direct limit of $G_0 \xrightarrow{\phi_0} G_1 \xrightarrow{\phi_1} G_2 \xrightarrow{\phi_2} \dots$.*

Proof. Obviously, G is isomorphic to the direct limit of $G_0 \xrightarrow{\phi_0} G_1 \xrightarrow{\phi_1} G_2 \xrightarrow{\phi_2} \dots$. The groups $G_0, G_1, G_2, \dots$ are hyperbolic by the Corollary 5.3.1 of Proposition 5.3. $\square$

It is easy to see that every recursively presented group with a finite regular file basis has solvable word problem. (Note that if a finitely generated group has a recursive presentation with a countably infinite alphabet, then it also has a recursive presentation with a finite alphabet.) In particular, if the group G is recursively presented, then it has solvable word problem. Generally, a recursively presented finitely generated group H has solvable word problem if and only if there exists a recursive set of group words X in a finite alphabet $\mathfrak{B}$ such that relative to some mapping of $\mathfrak{B}$ to H , every element of H has a unique representative in X . In the case of G , the set of all regular group words over $\mathfrak{A}$ may be taken as such a recursive set of representatives.

Property 5 (conditional). *If G is recursively presented, then it has solvable conjugacy problem.*

Proof. Assume G is recursively presented. Then, as noted above, G has solvable word problem.

Let q be an integer such that $q \geq 1/(1 - 2\mu)$.

Here is a description of an algorithm that solves the conjugacy problem in G :

Input: group words u and v (in the alphabet $\mathfrak{A}$).

Step 1: If $[u] \neq 1$ and $[v] \neq 1$, go to the next step; otherwise, test whether $[u] = [v]$. If yes, output ‘‘Yes’’, if no, output ‘‘No’’. Stop.

Step 2: Produce the (finite) set $\mathcal{S}$ of all group words s over $\mathfrak{A}$ such that $[s] = 1$ and $|s| \leq q(|v| + |u|)$.

Step 3: Produce a (finite) set D of up to isomorphism all annular diagrams over $\langle \mathfrak{A} \parallel \mathcal{S} \rangle$ with up to $q(|v| + |u|)$ edges.

Step 4: If there is $\Delta \in D$ such that u is the label of a representative of one of the contours of Δ and v^{-1} is the label of some representative of the other, then output ‘‘Yes’’; otherwise, output ‘‘No’’. Stop.

On an input (u, v) , this algorithm gives ‘‘Yes’’ as the output if and only if $[u]$ and $[v]$ are conjugate in G ; otherwise, it gives ‘‘No.’’ Below is an outline of a proof.

It is clear that the algorithm gives ‘‘Yes’’ only if $[u]$ and $[v]$ are conjugate in G . It is also clear that on every input it gives either ‘‘Yes’’ or ‘‘No,’’ and terminates.

Now, let u and v be arbitrary group words representing conjugate elements of G , and consider (u, v) as an input to the algorithm.

Case 1: $[u] = 1$ or $[v] = 1$. Then $[u] = 1 = [v]$, and therefore the algorithm gives “Yes” on Step 1.

Case 2: $[u] \neq 1$ and $[v] \neq 1$. Let Δ be a weakly reduced annular disc diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ such that u and v^{-1} are the labels of some representatives of its two contours (such Δ exists by Lemma V.5.2 in [LS01]). Let $\mathcal{S}$ be the set of group words produced by the algorithm on Step 2. Every element of $\mathcal{R}$ of length at most $q(|v| + |u|)$ is in $\mathcal{S}$. Let D be the set of annular diagrams produced by the algorithm on Step 3. By Proposition 5.1 and the Main Lemma,

$$|u| + |v| \geq (1 - 2\mu) \sum_{\Pi \in \Delta(2)} |\partial \Pi| \geq \frac{1}{q} \sum_{\Pi \in \Delta(2)} |\partial \Pi|$$

(recall that a special selection exists on every diagram over $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$). Therefore, the length of the contour of every face of Δ is at most $q(|u| + |v|)$, and the total number of edges of Δ is

$$\begin{aligned} \|\Delta(1)\| &= \frac{1}{2} \left(\sum_{\Pi \in \Delta(2)} |\partial \Pi| + |u| + |v| \right) \leq \frac{1}{2} (q(|u| + |v|) + |u| + |v|) \\ &= \frac{q+1}{2} (|u| + |v|) \leq q(|u| + |v|). \end{aligned}$$

Hence, Δ is isomorphic to some element of D , and the algorithm returns “Yes” on Step 4. $\square$

Proof of the theorem (see the Introduction). Let N be an integer such that $\lambda_1 n N \geq 1$. Let the construction in Section 2 be carried out under the following two additional conditions:

- (1) the order imposed on the set of all reduced group words is deg-lex;
- (2) for every i for which r_i is defined, $m_i = N|w_i| + i$.

Evidently, these conditions are consistent with the rest.

By Proposition 5.4, the group G is recursively presented. Properties 1–5 show that up to isomorphism G is a desired group. (G is *isomorphic* to a direct limit of a sequence of hyperbolic groups with respect to a family of surjective homomorphisms, but a desired group must *be* such a limit.) $\square$

6. COMMENTS

The group G and its presentation $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ mentioned in this section are defined in Section 2.

Proposition 6.1. *No hyperbolic group can be used as an example to demonstrate the negative answer to Bludov’s question.*

Proof. Suppose a hyperbolic group H is an example demonstrating the negative answer. Then H is *boundedly generated* (is the product of a finite sequence of its cyclic subgroups) and not virtually polycyclic. In particular,

H is non-elementary. Corollary 4.3 in [Min04] states that every boundedly generated hyperbolic group is elementary. (This fact also follows from Corollary 2 in [Ol'93].) This gives a contradiction. $\square$

Corollary 6.1.1. *The group G is not finitely presented.*

Proof. Suppose G is finitely presented. Then the presentation $\langle \mathfrak{A} \parallel \mathcal{R} \rangle$ is finite. By Corollary 5.3.1 of Proposition 5.3, G is hyperbolic. This contradicts Proposition 6.1. $\square$

Thus, the set $\mathcal{R}$ and the sequences $\{w_i\}_{i=1,\dots}$, $\{m_i\}_{i=1,\dots}$, $\{r_i\}_{i=1,\dots}$ constructed in Section 2 are infinite.

ACKNOWLEDGEMENTS

The author thanks Alexander Ol'shanskiy for suggesting the problem and engaging in fruitful discussions during the course of this work.

REFERENCES

- [ABC⁺91] Juan M. Alonso, Thomas Brady, Daryl Cooper, Vincent Ferlini, Martin Lustig, Michael L. Mihalik, Michael Shapiro, and Hamish B. Short, *Notes on word hyperbolic groups*, Group theory from a geometrical viewpoint, World Sci. Publ., River Edge, NJ, 1991, Proceedings of the workshop held in Trieste, March 26 – April 6, 1990, pp. 3–63.
- [Blu95a] Vasilij Vasilievich Bludov, *Nitevye bazisy v gruppakh [File bases in groups]*, Algebra i logika **34** (1995), no. 3, 247–261, in Russian.
- [Blu95b] ———, *File bases in groups*, Algebra and Logic **34** (1995), no. 3, 131–139, translated from Russian.
- [Bro94] Kenneth S. Brown, *Cohomology of groups*, Springer-Verlag, 1994, corrected reprint of the 1982 original.
- [CCH81] Ian M. Chiswell, Donald J. Collins, and Johannes Huebschmann, *Aspherical group presentations*, Math Z. **178** (1981), no. 1, 1–36.
- [Hue79] Johannes Huebschmann, *Cohomology theory of aspherical groups and of small cancellation groups*, J. Pure Appl. Algebra **14** (1979), no. 2, 137–143.
- [Hue80] ———, *The homotopy type of a combinatorially aspherical presentation*, Math. Z. **173** (1980), no. 2, 163–169.
- [Itô55] Noboru Itô, *Über das Produkt von zwei abelschen Gruppen*, Math. Z. **62** (1955), 400–401, in German.
- [LR80] John C. Lennox and James E. Roseblade, *Soluble products of polycyclic groups*, Math. Z. **170** (1980), no. 2, 153–154.
- [LS01] Roger C. Lyndon and Paul E. Schupp, *Combinatorial group theory*, Springer-Verlag, 2001, Reprint of the 1977 edition.
- [Min04] Ashot Minasyan, *On products of quasiconvex subgroups in hyperbolic groups*, Internat. J. Algebra Comput. **14** (2004), no. 2, 173–195.
- [MK95a] V. D. Mazurov and E. I. Khukhro (eds.), *Kourovskaja tetrad': nereshennyye voprosy teorii grupp [The Kourovka Notebook]*, 13th augmented ed., Inst. Mat. SO RAN, Novosibirsk, 1995, in Russian.
- [MK95b] V. D. Mazurov and E. I. Khukhro (eds.), *The Kourovka Notebook: unsolved problems in group theory*, 13th augmented ed., Russian Acad. of Sci. Siber. Div., Inst. Math., Novosibirsk, 1995, translated from Russian.
- [Mur05] Alexey Yu. Muranov, *Diagrams with selection and method for constructing boundedly generated and boundedly simple groups*, Comm. Algebra **33** (2005), no. 4, 1217–1258, arXiv.org preprint: math.GR/0404472.
- [Ol'89] Alexander Yu. Ol'shanskii, *Geometrija opredelajushchikh sootnoshenij v gruppakh [Geometry of defining relations in groups]*, Nauka, Moscow, 1989, in Russian.

- [Ol'91] ———, *Geometry of defining relations in groups*, Kluwer Academic Publishers, Dordrecht, Boston, 1991, translated from Russian.
- [Ol'93] ———, *On residualizing homomorphisms and G -subgroups of hyperbolic groups*, Internat. J. Algebra Comput. **3** (1993), no. 4, 365–409.

DEPT OF MATHEMATICS, VANDERBILT UNIVERSITY, 1326 STEVENSON CENTER, NASHVILLE, TN 37240–0001, USA.

Current address: Institut Camille Jordan, Université Lyon 1, 43 blvd du 11 novembre 1918, 69622 Villeurbanne cedex, France.

E-mail address: `muranov@math.univ-lyon1.fr`